## Minkowski's foundations of spacetime physics and **SDSS data motivate reassessment of ACDM cosmology**

"Failing to grasp the extent of the subversion of academia causes" them to try hard to force fit their discoveries into false models."

**Alexander F. Mayer** 

SensibleUniverse.net

-Asha Logos



The objective, statistical nature of SDSS astrophysical datasets, which were not driven by any theoretical agenda, reveal false and misleading prior measurements (e.g., redshift-distance) driven by confirmation bias in the context of such agendas. SDSS theta-z, redshift-magnitude (both spectroscopic and photometric pipelines), and galaxy population-density data are shown to conflict with the ACDM standard cosmological model. However, all four of those distinct and independent data sets are similarly consistent with a new cosmological model that revives de Sitter's 1917 solution to the field equations, long thought to entail an "empty universe." That new model, which represents a paradigm shift in cosmology, derives from considerations of symmetry and local proper time modeled as a geometric object, motivated by Minkowski (1909). The confrontation of all new predictive equations with corresponding SDSS data sets, using no free parameters, definitively resolves the modern quandary of astrophysical observables interpreted as accelerating cosmic expansion induced by 'dark energy.' The canonical idea of a non-relativistic universal time coordinate (i.e., ~13.7 Gyr of 'Cosmic Time' from initial singularity) is supplanted by a relativistic, strictly-local time coordinate involving no such inscrutable singularity.

## Abstract





The talk on "Space and Time," which Hermann Minkowski gave at the Convention of German Scientists and Doctors in Cologne, is the last of his ingenious creations. Unfortunately, it was not destined for him *to finish the more detailed development of his audacious concept* of a mechanics in which time is integrated with the three dimensions of space.

Since my student years Minkowski was my best, most dependable friend who supported me with all the depth and loyalty that was so characteristic of him. ...what death cannot take away is his noble image in our hearts and the knowledge that his spirit in us continues to be active.

# - A. Gutzmer

David Hilbert

You are encouraged to review this fact-finding and independently reproduce all analyses and results. Due to the unusually disruptive nature of this work, scientific integrity must overrule personal agendas.

Online self-study slides\* including copious text are obviously not intended for audience presentation; this PDF is a digital research report and textbook.

\* Internet links are attached to references, names, and graphics in the PDF; cursor typically switches to a hand pointer when hovering over the links. A minority of all available links are highlighted with the link  $N_{O}$  indicator.



# Sloan Digital Sky Survey

This presentation is © 2018–2019 A. F. Mayer.

Image credit: Reidar Hahn, Fermilab Visual Media Services



**Sloan Foundation** 2.5-m Telescope Apache Point, NM



# **Digital Research Report and Instructional Textbook**

- Table of Contents – SDSS THETA-Z DATA 15 SDSS REDSHIFT-MAGNITUDE DATA 57 – DE SITTER COSMOLOGY REVISITED 99 – NEW PREDICTIVE EQUATIONS 130 GALAXY SPACE DENSITY 142 **CREDITS AND AFTERWORD** 150 ADDENDUM 1: RELATED COSMOLOGICAL ISSUES 161 – ADDENDUM 2: RELEVANT HISTORY, ETC. 187
- PROLOGUE INSTRUMENTS AND TOOLS PARTI PART II PART III PART IV PART V PART VI PART VII PART VIII



## PROLOGUE – INSTRUMENTS AND TOOLS

Apache Point Observatory Sunspot, New Mexico, USA



Ultraviolet	Green	Red	Near Infrared	Infrared	
u	g	r	i	Z	
3531	4627	6140	7467	8887	




photo credit: Smithsonian National Air and Space Museum

SDSS astronomical filters  $\lambda_{eff}$  (Å)

Herein, not to be confused with *redshift* (z)!

Central wavelengths  $\lambda_{eff}$  (Å) from Doi et al. (2010), Table 2.






photo credit: Michael Carr







#### **NEW: SciServer Betelgeuse (v2.0.0)!**

We are proud to announce a new release of the SciServer online science platform! The "Betelgeuse" release has many new features to manage and share big data resources. about the changes on the SciServer website!

#### Welcome to the DR14 site!!!

This website presents data from the Sloan Digital Sky Survey, a project to make a map of a large part of the universe. We would like to show you the beauty of the universe, and share with you our excitement as we build the largest map in the history of the world.

#### News

The site hosts data from D Release 14 (DR14). What DR14, and known proble More...

#### **Data Access**

**Finding Chart** Navigate Quick Look | Explore Image List Search IQS | SQS | IRSQS SQL Search Cross-ID SkyQuery CrossMatch NEW! CasJobs 🛛 <

#### Education


For Educators Lesson Plans **College Lab Activities** Instructor Guides Student/Public Research Galaxy Zoo 🗾 Zooniverse 🗖 Voyages 🗖

#### Links

sdss.org 🗗 Data Release 14 🕑 Surveys 🖓 Instruments 🦉 SDSS Science Science Archive Server About Astronomy About SkyServer Credits 🗖

#### Help

Start Here Glossary Algorithms 岱 Cooking with SQL Tutorial About the Data Schema Brows Sample SQL Q







**Contact Us** 



skyserver.sdss.org/dr14/en/home.aspx


# SciServer (

Site Search	n Help	History NEW!	afmayer
e new . Learn	SDSS is supported by		URL skyserver.sdss.org
	SSF C		
Data at's new in ems.	Powered by		
FAQ 🖾	SciServer		
<u></u>	Microsoft		
Sloan	Site Traffic Privacy Policy		
tabase			
rser Queries			Catalog Archive Server (CAS)
a different regions of a Xiplane, interneting the one bill another than a second second and the programmers of a Xib hortzet water instructions reach and travellater, and a more than second reaches and travellater,			

C




SDSS	Quer <mark>Query</mark>		sJobs			
Help Tools Query	History	MyDB	Import	Groups	Output	Sc
Context DR14 Samples Recent Clear	default	ch Table	(optional) R50_g	Task Nar		[ ∕∕ly
1 SELECT S.Z S.Z S.ZErr , g.petroR50 , g.petroR50E FROM Galaxy g SpecObj s 9 WHERE	rr_g				Change to get Change	da <sup>-</sup> "R!
10 s.programna 11 AND s.zWarning 12 AND s.zErr >= 0 13 AND ABS(s.zErr/ 14 AND s.z > 0 15 AND g.clean = 1 16 AND (g.flags_g 17 AND (g.flags_g 18 AND g.petroR50 19 AND g.petroR50 20 AND g.petroR50 21 AND g.specObjID	= 0 s.z) <= & 0x100 & 0x200 g > 0 rr_g >= rr_g /	0.01 = 0 = 0 0 g.petrol	R50_g <	0.20	disallo disallo disallo disallo disallo disallo disallo disallo disallo max. mo join to	ow edsl ow pho ow ow ow ow




to "\_" with r , i , z for those filters. 50" to "Rad" for that data. **Run SQL statements** on "DR14" database (Data Release 14).

minor surveys redshifts with warnings -n error codes hift error is 1% [÷ 0], above tometry NOPETRÒ flag MANYPETRO flag -9999 error code -1000 error code urement error is 20% les on specObjID key

SDSS data releases are cumulative. DR 15 (10 Dec. 2018) includes new, additional data that should not alter the prior SDSS data (DR 14) and its analysis that are presented herein.




	skyserver.sdss.org/CasJobs/MyDB.aspx	0	
	OB tab	Outputer SkySoryer	SciServer afmayer
MyDB    Local Only    Views Tables Functions Procedures			
Sort by Rows kB Name 806,483 20,040 petroR50_g (1) Click on		petroR50Err_g real [4]	
the filename	Notes No notes found		
	New Note		



# Free, cross-platform astrophysical plotting package

## http://www.star.bris.ac.uk/~mbt/topcat/



## **Tool for OPerations on Catalogues And Tables**

Does what you want with tables

Latest (see <u>Version History</u> for details)

Version 4.6-3 released 9 May 2019

#### **TOPCAT** Creative



Dr. Mark Taylor Ph.D. (Bristol)

Astrophysics group University of Bristol






Screenshots Help & Support Development News Home Examples Download Features

#### Veusz – a scientific plotting package

Veusz is a scientific plotting and graphing program with a graphical user interface, designed to produce publication-ready 2D and 3D plots. In addition it can be used as a module in Python for plotting. Veusz is multiplatform, running on Windows, Linux/Unix and macOS. It supports vector and bitmap output, including PDF, Postscript, SVG and EMF. Veusz is Free Software.

With the help of a tutorial the program can be used by the novice user and is flexible for advanced work. In Veusz plots are created by building up plotting widgets with a consistent object-based interface, where the user sets the properties of the widgets. There are many options for customization of plots. See features, 2D examples and 3D examples to see what can be done with the program.

It allows data to be imported from text, CSV, HDF5 and FITS files. Datasets can also be entered within the program and new datasets can be created via the manipulation of existing datasets using mathematical expressions and more. The program can also be extended, by adding plugins supporting importing new data formats, different types of data manipulation or for automating tasks.



# Free and cross-platform

Ċ

#### https://veusz.github.io



# PART I – SDSS THETA-Z DATA


Note: slides 16–28 are a precursor to analysis.

Click to go back to Table of Contents...

# Understanding SDSS angular resolution ("theta" measurement) The *average* apparent diameter of the full Moon is 31.1' = 1866''.



The midline shown has a width of 10" (i.e., 10 arcseconds), and the green dot at 6 o'clock position has a 10" diameter, as compared to the average apparent size of the full Moon.



## SDSS glossary reference

#### petroRad


The Petrosian radius. A measure of the angular size of an image, most meaningful for galaxies. Units are seconds of arc. The Petrosian radius (and related measures of size called petroR50 and petroR90) are derived from the surface brightness profile of the galaxy, as described in Algorithms.

#### **Surface Brightness**

The frames pipeline also reports the radii containing 50% and 90% of the Petrosian flux for each band, petroR50 and petroR90 respectively.



## Petrosian radius




Here, r is a variable measurement; it is not a recorded measurement.


Define  $\mathscr{R}_{P,lim}$  as  $\mathscr{R}_{P}(r_{P}) = \mathscr{R}_{P,lim}$  where  $r_{P}$  is the measured and recorded Petrosian radius.  $\mathscr{R}_{P,lim} = 0.2$  for the SDSS:

Varying r, when  $\mathscr{R}_{P}(r) = 0.2$ , then  $r_{P} = r$ .

**Reference:** https://www.sdss.org/dr14/algorithms/magnitudes/#mag\_petro

#### DEFINITION: "Petrosian ratio," $\mathscr{R}_{P}(r) =$






local surface brightness in annulus (blue region) mean surface brightness measured inside radius r


$$\mathscr{R}_{P}(r) \equiv \frac{\int_{0.8r}^{1.25r} dr' 2\pi r' I(r') / \pi (1.25^{2} - 0.8^{2}) r^{2}}{\int_{0}^{r} dr' 2\pi r' I(r') / \pi r^{2}}$$

•


The observed radial galaxy brightness profile [I(r')] is "azimuthally averaged" (i.e., averaged over  $2\pi$  radians).








## $\sim 804 \times 10^3 \; \texttt{galaxies}$ plotted






# $\sim 1.372 \times 10^{6} \; \texttt{galaxies}$ plotted


# SpecObj.z (redshift)


## Theta-z (half-light radius)



# $\sim 1.498 \times 10^{6} \ {\rm galaxies} \ {\rm plotted}$

# SpecObj.z (redshift)







# $\sim 966 \times 10^3 \; \texttt{galaxies}$ plotted Avg. error: 8.5%


0.1 0.2 SpecObj.z (redshift)

0.5






0.5 0.1 0.2 SpecObj.z (redshift)









# $\sim 719 \times 10^3 \; \texttt{galaxies}$ plotted

24

SpecObj.z (redshift)

## **Empirical inference**

The *intrinsic* radial size of *typical* galaxies, including all morphologies, varies by about one order of magnitude (i.e., x10). That decadal range excludes the statistically-unlikely outliers.

## SQL query for theta-z data...

```
SELECT
    S.Z
    s.zErr
,
    g.petroR50_g
    g.petroR50Err_g
FROM
    Galaxy g
    SpecObj s
WHERE
    s.programname IN ('legacy', 'boss') -- disallow minor surveys
AND s.zWarning = 0
AND s.zErr >= 0
AND ABS(s.zErr/s.z) \leq 0.01
AND s.z > \emptyset
AND g.clean = 1
AND (g.flags_g & 0 \times 100) = 0
AND (g.flags_g & 0 \times 200) = 0
AND g.petroR50_g > \emptyset
AND g.petroR50Err_g >= 0
AND g.petroR50Err_g / g.petroR50_g < 0.20 -- max. measurement error is 20%
AND g.specObjID = s.specObjID
```

- -- Change "\_g" to "\_" with r , i , z
- -- to get data for those filters.
- -- Change "R50" to "Rad" for that data.

- -- disallow redshifts with warnings
- -- disallow -n error codes
- -- max. redshift error is 1%
- -- disallow [÷ 0], above
- -- clean photometry
- -- disallow NOPETRO flag
- -- disallow MANYPETRO flag
- -- disallow -9999 error code
- -- disallow -1000 error code
- -- join tables on specObjID key

# SQL modifications for \_gri (average of {g, r, i})

#### SELECT

. . .

S.Z

```
s.zErr
,
```

- (g.petroR50\_g + g.petroR50\_r + g.petroR50\_i) / 3 AS petroR50\_gri

## Add to WHERE clause:

- AND (g.flags\_r &  $0 \times 100$ ) = 0
- AND  $(g.flags_r \& 0 \times 200) = 0$
- AND g.petroR50\_r > 0
- AND g.petroR50Err\_r >= 0
- AND g.petroR50Err\_r / g.petroR50\_r < 0.20 -- max. measurement error is 20%

AND (g.flags\_i &  $0 \times 100$ ) = 0 AND  $(g.flags_i \& 0 \times 200) = 0$ AND g.petroR50\_i > 0 AND g.petroR50Err\_i >= 0

(g.petroR50Err\_g + g.petroR50Err\_r + g.petroR50Err\_i) / 3 AS petroR50Err\_gri

- -- disallow NOPETRO flag
- -- disallow MANYPETRO flag
- -- disallow -9999 error code
- -- disallow -1000 error code
  - -- disallow NOPETRO flag
  - -- disallow MANYPETRO flag
  - -- disallow -9999 error code
  - -- disallow -1000 error code
- AND g.petroR50Err\_i / g.petroR50\_i < 0.20 -- max. measurement error is 20%

## SQL modifications for average error statistic

#### SELECT AVG(petroR50Err\_z / petroR50\_z) \* 100 AS AVGpetroR50Err\_z


. . .

. . .

-- When averaging the three filters {g, r, i} use: SELECT AVG((petroR50Err\_g / petroR50\_g + petroR50Err\_r / petroR50\_r + petroR50Err\_i / petroR50\_i) / 3) \* 100 AS AVGpetroR50Err\_gri









#### Redshift bins (0.02, 0.04, 0.08, 0.16, 0.32, 0.64)

SpecObj.z (redshift)







## $z \approx 0.04$

## $(0.038 \le z \le 0.042) \& n > 6$

Recall that in a Euclidean space, the doubling z apparent size of an object is inversely each time... proportional to distance (i.e., half size at double the distance). .34″  $\mathbf{C}$ nean: 10 5 Galaxy.petroR50\_gri (arcsec)











#### $z \approx 0.08$


## $(0.076 \le z \le 0.084) \& n > 6$











## SQL for redshift-bin histograms

```
SELECT
, COUNT(1) AS n
```


# Append to \_gri WHERE clause:

. . .

AND S.Z BETWEEN 0.018 AND 0.022 -- CHANGE THIS FOR EACH BIN. **GROUP** BY  $ROUND((petroR50_g + petroR50_r + petroR50_i) / 3, 1)$ HAVING COUNT(1) > 3 -- CHANGE THIS AS NEEDED. ORDER BY 1

#### ROUND((petroR50\_g + petroR50\_r + petroR50\_i) / 3 , 1) AS petroR50\_gri

#### Mean measurement of Galaxy.petroR50\_gri within each redshift bin



SpecObj.z (redshift)

37

White line is a visual aid.

## Quantitative analysis of galaxy-size statistical distribution (z = 0.08)

#### SELECT ROUND(AVG(z), 3) AS AVGz , , , , COUNT(1) AS n

# Append to \_\_gri WHERE clause:

. . .


AND s.z BETWEEN 0.076 AND 0.084 -- z = 0.08 bin

AVGz	ArithMean	StdDev	mu	sigma	n
0.080	2.45	1.01	0.8194	0.3886	35985

 $ROUND(AVG((petroR50_g + petroR50_r + petroR50_i) / 3), 2) AS ArithMean$ ROUND(STDDEV((petroR50\_g + petroR50\_r + petroR50\_i) / 3), 2) AS StdDev  $ROUND(AVG(LN((petroR50_g + petroR50_r + petroR50_i) / 3)), 4) AS mu$ ROUND(STDDEV(LN((petroR50\_g + petroR50\_r + petroR50\_i) / 3)), 4) AS sigma

## Empirical distribution vs. log-normal probability density function (PDF)


2 k  $\bar{z} = 0.080$  $(0.076 \le z \le 0.084)$ 35,985 galaxies 57 u )  $\mathbf{H}$ P  $(\ln heta - \mu)$  $2\sigma^2$ **PDF** count  $\theta \sigma \sqrt{2\pi}$ 1K Bin 500 0 0.2 0.5




39

Galaxy.petroR50 gri (arcsec)

# Quantitative analysis of galaxy-size statistical distribution (all z-bins)



## QQ plot confirmation of log-normal galaxy-size distribution



#### SELECT (petroR50\_g + petroR50\_r + petroR50\_i) / 3), 2) AS petroR50\_gri

# Append to \_\_gri WHERE clause:

. . .

-- Uncomment one line, below, to choose that bin for analysis. -- AND s.z BETWEEN 0.018 AND 0.022 -- z = 0.02 bin -- AND s.z BETWEEN 0.038 AND 0.042 -- z = 0.04 bin AND s.z BETWEEN 0.076 AND 0.084 - z = 0.08 bin -- AND s.z BETWEEN 0.152 AND 0.168 -- z = 0.16 bin -- AND s.z BETWEEN 0.304 AND 0.336 -- z = 0.32 bin -- AND s.z BETWEEN 0.608 AND 0.672 -- z = 0.64 bin

# SQL for QQ plot

# Quantitative analysis of galaxy-size statistical distribution (all z-bins)...

AVGz	ArithMean	StdDev	mu	sigma	n
0.020	4.75	2.83	1.4014	0.5627	3186
0.040	3.41	1.68	1.1178	0.4688	7834
0.080	2.45	1.01	0.8194	0.3886	35985
0.160	2.13	0.66	0.7114	0.2994	32942
0.320	1.60	0.39	0.4436	0.2367	34769
0.636	1.20	0.28	0.1533	0.2278	7036

# **Empirical inference**

# Galaxies have a log-normal size distribution, which is clearly observable over a wide range of redshift.



# **Empirical inference**

To *good approximation*, the mean apparent galaxy size in each redshift bin can be expected to represent the same standard rod (i.e., to be very nearly the same *intrinsic* size). To assume otherwise would require some unphysical ad hoc explanation.

With a sample population across six redshift bins of ~122k galaxies, an increase in redshift of  $32 \times (z: 0.02 \rightarrow 0.64)$  is correlated to a smooth and continuous decrease in mean apparent half-light radius of about  $4 \times$ . This is eight times less than expected as per the 'Hubble law' predictive model.

3


"And where science gets *tough*, tough in the sense that you can be wrong, even if you passionately believe something, is that a good scientific theory makes predictions and those predictions can be tested."

Online Lecture: The Nature of Space and Time (1:05:50 / 1:28:25)

- Lee Smolin, Perimeter Institute for Theoretical Physics







Bin redshift (z)

#### Let's be clear about what that graph means, as it is non-trivial:


First, we confirm that the empirical SDSS theta-z data is objective and that there is no systematic error or any other problem that invalidates logical inferences that may be drawn from that data.

The graph means that the empirical redshift-distance relationship is inconsistent with the 'Hubble law.' If that is true, then the Universe is not expanding, regardless of what we may *believe* or want to believe. The data is unambiguous; it is based on  $\sim 2 \times 10^6$  objective high-quality measurements that are not affected by confirmation bias.

Accordingly, we have a major *scientific crisis* (confirmed by several other independent, corroborating empirical observations, shown later). To find a resolution, there is only one place to look for guidance...


# The Einstein Field Equations

(The physical constants are normalized.)





# Willem de Sitter (6 May 1872 – 20 Nov 1934) was a Dutch mathematician, physicist, and astronomer.



publications link dwc.knaw.nl ⇒



#### uc KNAW Digital Library




In a nutshell, we go back to Willem de Sitter's exact solution to the Einstein field equations (EFE) circa 1917 and gain a clear and rational understanding of its *physical interpretation*, which new interpretation supersedes that of the Einstein-de Sitter debate.<sup>1</sup> This leads to the derivation of a new set of predictive formulas, which rest on first principles and are consistent with the EFE. Among these formulas is:


$$\theta(z) = C_{\delta} \left( 1 - \frac{1}{(z+1)^2} \right)^{-\frac{1}{2}} \text{ radians}$$

arXiv:1503.08029 [physics.hist-ph].

 $C_{\delta}$  is an arbitrary scaling constant that is proportional to the intrinsic diameter ( $\delta$ ) of the astrophysical standard rod under consideration. Thus, changing  $C_{\delta}$  shifts the predictive curve up or down to match an arbitrary object size, but it does not modify that curve in any way: With regard to relative measurements, there are no free parameters in any of the new predictive formulas. For absolute measurements, the single shared free parameter is the estimated Cosmic radius R.

1. C. O'Raifeartaigh, M. O'Keeffe, W. Nahmb and S. Mitton, "Einstein's cosmology review of 1933: a new perspective on the Einstein-de Sitter model of the cosmos," Euro. Phys. J. H. 40, 301 (2015); 5 I





- Bin redshift (z)


### Let's be clear about what the red curves on the prior two slides mean, as it is non-trivial:

As an accurate predictive model of empirical observations, they mean that the Universe is not expanding, regardless of what we may *believe* or want to believe. Over a century ago, de Sitter *unequivocally* pursued the right idea (galaxy redshifts are caused by cosmological spacetime geometry and not by expansion-induced recession). However, he did not adequately interpret (and further develop) his own mathematics.

### Quoted from Edwin Hubble's last paragraph; PNAS 15, 168 (1929)


P

The outstanding feature, however, is the possibility that the velocitydistance relation may represent the de Sitter effect, and hence that numerical data may be introduced into discussions of the general curvature of space. ... it may be emphasized that the linear relation found in the present discussion is a first approximation representing a restricted range in distance.





# 1929 Hubble diagram (annotated)





# PART II – SDSS REDSHIFT-MAGNITUDE DATA

Click to go back to Table of Contents...

$$m = -2.5 \cdot \log_{10}$$

- *Brighter* objects have a *smaller* magnitude.
- -5 "mags" is exactly 100× brighter.
- $-1 \text{ mag is } \sqrt[5]{100} \approx 2.512 \times \text{ brighter.}$
- $\Delta m$  mags is a change in brightness of  $2.512^{-\Delta m} \times$

- The astronomical magnitude scale
- Apparent magnitude is quantified in terms of the radiation *flux ratio* within a given spectral band (x):




# Three among the typical, smaller galaxies in the redshift bin (petroR50 gri=1.8'')

petroR50\_gri ( $\pm 0.1''$ )




### $z = 0.080 \pm 0.004 \ (\pm 5\%)$



Click images for SDSS SkyServer Explorer.

### ○ 3"-diameter SDSS fiber

## ○ 2"-diameter SDSS fiber






# Three among the largest galaxies in the redshift bin, including the extreme outlier

petroR50\_gri ( $\pm 0.1''$ )



## $z = 0.080 \pm 0.004 (\pm 5\%)$



*Change of scale* (demagnified 2×) → SCALE: 0.0853 arcsec/pixel


Click images for SDSS SkyServer Explorer.

#### ○ 3"-diameter SDSS fiber

# ○ 2"-diameter SDSS fiber







# **Empirical inference**

By inspection\* : to close approximation, the intrinsic diameter of galactic nuclei are constant and independent of host-galaxy size.

Also, down to about  $z \sim 0.08$ , a 3"-diameter fixed aperture encloses the apparent diameter of galactic nuclei, thus measuring their flux. Any distal regions of more distant galaxies so enclosed will add a much smaller contribution to the total flux measured by the fiber.

\* (of numerous randomly-selected SDSS galaxy images)

(4)

### Fixed-aperture fiber magnitudes


These data are important because they are *independent* of the photometricpipeline magnitude measurements (e.g., Petrosian magnitudes), with which they shall be shown to be in agreement as concerns empirical interpretation.

Fixed-aperture fiber magnitudes are a *byproduct* of the optical spectroscopy measurements (e.g., redshift)—as such, they were not previously considered to be photometric "science data." However, by serendipitous correlation, the 3"- and 2"-diameter (Legacy and BOSS spectrograph fibers, respectively) measure (to good approximation over  $0.08 \le z \le 0.5$ ) the apparent brightness of galactic nuclei, which dominate the total light flux of galaxies. Just like stars, these ubiquitous objects have a span of intrinsic brightness; given data on  $\sim 10^6$  objects, *population statistics* can be exploited for scientific discovery.

# Fixed-aperture effect over a wide range of redshift (0.02, 0.08, 0.60)

3"-diameter SDSS fiber

# Aperture too small for nucleus:



Click images for SDSS SkyServer Explorer.



### 2"-diameter SDSS fiber

# Aperture encloses nucleus:

# Aperture too large for nucleus:

63

#### SCALE: 0.0427 arcsec/pixel

## SDSS documentation reference

### Fiber Magnitudes: fiberMag and fiber2Mag

Fiber magnitudes reflect the flux contained within the aperture of a spectroscopic fiber in each band. In the case of fiberMag we assume an aperture appropriate to the SDSS spectrograph (3" in diameter). In the case of fiber2Mag we assume an aperture appropriate to the BOSS spectrograph (2" in diameter).

#### Notes

- $\bigcirc$
- $\bigcirc$ the fiber magnitudes closer to what is seen by the spectrograph.
- $\bigcirc$

For children of deblended galaxies, some of the pixels within a 1.5" radius may belong to other children; we now measure the flux of the parent at the position of the child; this properly reflects the amount of light which the spectrograph will see.

Images are convolved to 2" seeing before fiberMags are measured. This also makes

Addendum to the above from DR7 notes: There is a cut in 3'' fiber magnitude at g=15, r = 15, and i = 14.5 (to prevent saturation and cross-talk in the spectrographs).<sup>1</sup>



# SDSS glossary reference

Fiber

The SDSS spectrograph uses optical fibers to direct the light at the focal plane from individual objects to the slithead. Each object is assigned a corresponding fiberID. The fibers for SDSS-I/II were 3 arcsecs in diameter in the source plane; they are 2 arcsecs in diameter for BOSS. Each fiber is surrounded by a large sheath which prevents any pair of fibers from being placed closer than 55 arcsecs on the same plate (62 arcsecs for BOSS).

#### fiberMag

The magnitude measured by the frames pipeline to simulate the flux that would fall into a 3" fiber in typical seeing. Similarly, fiber2Mag simulates the 2" fiber magnitude.

#### Plate

Each spectroscopic exposure employs a large, thin, circular metal plate that positions optical fibers via holes drilled at the locations of the images in the telescope focal plane. These fibers then feed into the spectrographs. Each plate has a unique serial number, which is called *plate* in views such as SpecObj in the CAS.



Average apparent diameter of the full Moon (31.1')

NASA/IPAC Extragalactic Database data:

3

Blueshift: 0.001001 ± 0.000013 (-300 ± 4 km/s) *Effective redshift distance*: *z* << 0.001

#### Summary Statistics computed by NED from 199 Distance(s) in the literature:

Mean Metric Distance: 0.788 Mpc **Standard Deviation:** Minimum Distance: Maximum Distance:

0.163 Mpc 0.440 Mpc 2.800 Mpc

M31 image credit: The Sloan Digital Sky Survey Collaboration

#### SDSS telescope composite image of M31



#### 1 degree (60')

M32

Scale derived from 0.6087° M31–M110 angular separation.



M31 image credit: The Sloan Digital Sky Survey Collaboration

### Detailed perspective of M31

#### M32

15'

#### 0.25 degrees (15')



Viewed in the infrared (i.e., z-band) bright nuclei are considerably smaller than as viewed in the optical.

#### $\lambda \sim 24 \ \mu m$

M31 image credit: NASA / JPL-Caltech / K. Gordon (University of Arizona)

#### Spitzer satellite IR image of M31

M32

15

#### 0.25 degrees (15')



#### Similarly in the UV

FUV: (1344 – 1786),  $\lambda_{eff} = 1528$  (Å); white/blue NUV: (1771 – 2831),  $\lambda_{eff} = 2271$  (Å); orange M31 image credit: NASA/JPL-Caltech

#### GALEX satellite UV image of M31



#### 0.25 degrees (15')



#### Similarly in the UV



M31 image credit: NASA/Swift/Stefan Immler (GSFC) and Erin Grand (UMCP)

#### SWIFT satellite UV image of M31

M32


 $\bigcirc$ 

#### 0.25 degrees (15')






### Redshift-Magnitude (fiberMag\_z)




# $\sim 1.379 imes 10^{6} \; {\tt galaxies plotted}$




# $\sim 772 imes 10^3$ galaxies plotted

SpecPhoto.z (redshift)








#### $\sim 184 \times 10^3 \; \texttt{galaxies}$ plotted

### LEGACY LRG Survey subset (Luminous Red Galaxy)


0.5






#### $\sim 608 imes 10^3$ galaxies plotted

74



#### $\sim 220 imes 10^3$ galaxies plotted


75



### $\sim 48 imes 10^3$ galaxies plotted

76


### <code>Redshift-Magnitude</code> (fiberMag\_z) $\sim$ 19 imes 10<sup>3</sup> galaxies plotted




#### AGN subset

#### 0.05 0.1 0.2 SpecPhoto.z (redshift)

0.5





20

19

18

17

16

15

14

0.01

### 0.02

# 0.05

### $\sim 14 imes 10^3$ galaxies plotted


#### **BROADLINE** subset

0.1 0.2 SpecPhoto.z (redshift)


0.5









# $\sim 2.100 \times 10^{6} \; \texttt{galaxies}$ plotted Redshift-Magnitude (fiberMag\_z) Complete dataset with no measurement-error limits. 0.1 0.2 0.5 SpecPhoto.z (redshift)



#### $\sim 1.379 imes 10^{6} \; {\tt galaxies \; plotted}$ Redshift-Magnitude (fiberMag\_z)



80




#### Standard candles, redshift-magnitude, and K-correction

Interpreted in the context of the ACDM consensus cosmological model, the graph means that from 5.1 to 1.3 Gyr ago (z: 0.5  $\rightarrow$  0.1) the intrinsic brightness of galaxy nuclei, as measured by SDSS fiber magnitudes over this redshift range with good accuracy and consistency, decreased by a factor of about five (i.e.,  $0.2\times$ ) at an accelerating rate, and that this phenomenon has continued to date. Such an interpretation defies rational explanation; accordingly, one may conclude that this predictive redshift-magnitude model curve is inconsistent with the empirical data, which is to be expected given the radical failure of the correlated theta-z model.

An accurately-modeled redshift-magnitude curve represents the apparent luminosity of a standard candle over redshift, with an empirical cosmic standard candle being expected to exhibit imperfection in the form of a *narrow range* of intrinsic brightness. In turn, such a cosmic standard candle, observed through a particular bandpass filter, must trace the applicable K-correction curve for that filter.<sup>1</sup> The K-correction curve reflects redshift-induced observation of the typical variation in galaxy flux as a function of emission wavelength, which is graphed in the following slide...

1. I. V. Chilingarian, A-L Melchior, & I. Y. Zolotukhin, MNRAS 405, 1409 (2010).

### Galaxy spectrum demonstrating 4000-Å (Ca II) break







#### Follow the data...

According to conventional thinking in cosmology, the idea that a class of standard-candle galaxies might exist may seem to be a "wild hypothesis" that is inconsistent with what are assumed to be 'known facts,' so confusion and misunderstanding are to be expected. However, those problems can be resolved:

If a class of cosmic object is observed to trace the known K-correction curve over redshift, then, by definition, that class of object must be a standard candle. Confronted with such empirical data, one is not "hypothesizing" the existence of a galactic standard candle, one is *objectively observing* the phenomenon.


The statistical (fuzzy)<sup>1</sup> baseline of the SDSS redshift-magnitude measurements represents the class of brightest galaxies (Petrosian magnitudes for an entire galaxy or their nuclei as measured to good approximation by the spectrograph fiber magnitudes). If that baseline traces the K-correction curve, then we may infer with complete certainty that those objects are also a class of standard candle, without regard for any redshift-magnitude model.

SO, LET US FOLLOW THE RELIABLE SDSS DATA AND SEE WHERE IT LEADS...


1. R. Viertl, Statistical Methods for Fuzzy Data (Wiley, Chichester, 2011).

84

#### Redshift-Magnitude (fiberMag\_z)



#### Redshift-Magnitude (fiberMag\_i)






#### $\sim 2.032 imes 10^{6} \; {\tt galaxies plotted}$


86


#### Redshift-Magnitude (fiberMagr)





#### $\sim 1.843 imes 10^{6}$ galaxies plotted





A fundamental assumption associated with the current standard cosmological model is galaxy evolution over lookback time (i.e., galaxies observed at higher redshift are expected to be *intrinsically* younger). However, if that assumption is true, one should not see "red and dead" (i.e., intrinsically very old) high-redshift galaxies (1 < z < 2), yet we do.<sup>1</sup>


Observation of an *empirical* m(z) "K-correction" curve as seen in the last four slides is a dependable *empirical signature* of a galactic cosmic standard candle. Observation of the +1-mag 4000-Angstrom break, which initiates at  $z \sim 0.3$  in the g-band as illustrated in the prior slide, positively identifies the dataset baseline as a set of very-nearly-consistent cosmic 'standard candles' (i.e., having minor intrinsic-brightness variation in the set); no "hypothesis" is involved.


#### Note: $LCDM \equiv \Lambda CDM - Lambda Cold Dark Matter [consensus model]$

1. R. G. Abraham et al., Star-Forming, Recently Star-Forming, and "Red and Dead" Galaxies at 1 < Z < 2, in: R. De Grijs and R. M. González Delgado (eds), Starbursts, Astrophysics and Space Science Library, vol 329, (Springer, Dordrecht 2005).

### <code>Redshift-Magnitude</code> (fiberMag\_z) $\sim$ 1.379 imes 10<sup>6</sup> galaxies plotted

To close approximation, the data's statistical baseline for all four bandpass filters traces the K-correction curve, so it must represent standard candles (i.e., objects having similar *intrinsic* brightness to the same approximation). The intrinsic 'noise' around this baseline is less than  $\pm 0.2$  mag.





0.05 0.1 0.2 SpecPhoto.z (redshift)

24

23

22

21

20


19


18


17

"fuzzy" statistical baseline of data (±0.2 mag)

0.5







#### Redshift-Magnitude (fiberMag\_g)

#### $\sim 1.115 \times 10^{6} \; \text{galaxies plotted}$

### SDSS glossary reference

#### Magnitude, Petrosian


Stored as petroMag. For galaxy photometry, measuring flux is more difficult than for stars, because galaxies do not all have the same radial surface brightness profile, and have no sharp edges. In order to avoid biases, we wish to measure a constant fraction of the total light, independent of the position and distance of the object. To satisfy these requirements, the SDSS has adopted a modified form of the Petrosian (1976) system, measuring galaxy fluxes within a circular aperture whose radius is defined by the shape of the azimuthally averaged light profile. Details can be found in the Photometry section of the Algorithms pages and the Strauss et al. (2002) AJ paper on galaxy target selection. Model magnitudes share most of the advantages of Petrosian magnitudes, and have higher S/N; they are therefore used instead of Petrosian magnitudes for target selection in BOSS.

If you jumped here, click to go back to *Fixed-aperture fiber magnitudes*...





$$F_{\rm P} \equiv \int_{0}^{N_{\rm P}r_{\rm P}} {\rm d}r' 2\pi r' I(r') \quad \left[N_{\rm P} = 2.0\right]$$




The SDSS Petrosian flux in any band is defined as the flux within two Petrosian radii  $(2r_P)$ .

"In the SDSS five-band photometry, the aperture in all bands is set by the profile of the galaxy in the r band alone. This procedure ensures that the color measured by comparing the Petrosian flux  $F_{\rm P}$  in different bands is measured through a consistent aperture."


**Reference:** https://www.sdss.org/dr14/algorithms/magnitudes/#mag\_petro

**DEFINITION: "Petrosian flux"** 





SpecObj.z (redshift)



### $\sim 650 imes 10^3$ galaxies plotted

#### The 'alignment' of the LCDM curve with the data is *deceptive*: no Ca II break! Systematic error in petroRad\_g data has caused petroMag\_g adulteration.


mag

break

<sup>†</sup> Wright (2006, *PASP*, **118**, 1711)  $H_0 = 69.6$ ,  $\Omega_M = 0.286$ ,  $\Omega_\Lambda = 0.714$ 

File: https://archive.org/details/WrightLCDMCalcs

0.1 0.2 SpecObj.z (redshift)




### SQL query for all redshift-magnitude data

```
SELECT -- Change "_z" to "_" with i , r , g to get data for those filters.
    Ζ
   zErr
,
   fiberMag_z -- Change "fiber" to "petro" for that data.
,
   fiberMagErr_z
)
   CASE (legacy_target1 & 0x20) WHEN 0 THEN 0 ELSE 1 END AS fLRGsurvey
,
   programname
2
    subClass
FROM
    SpecPhoto
WHERE
    programname IN ('legacy', 'boss') -- disallow minor surveys
AND class = 'GALAXY''
AND zWarning = 0
AND zErr >= 0
AND ABS(zErr/z) <= 0.01
AND z > 0
ORDER BY
    Ζ
```


-- remove QSO and STARs -- disallow redshifts with warnings -- disallow -n error codes -- max. redshift error is 1% -- disallow [÷ 0], above AND fiberMagErr\_z BETWEEN 0 AND 0.1 -- disallow errors >0.1 mag and -9999 error code

#### 'de Sitter' model redshift-magnitude curve (constant intrinsic brightness)



### Redshift (z)






### PART III – DE SITTER COSMOLOGY REVISITED

Click to go back to Table of Contents...



### The Main Characters

4



Bernhard Riemann (1826 – 1866)

Lifetime list of 11 published papers



#### Albert Einstein (1879 – 1955)

List of scientific publications by Albert Einstein (Wikipedia)



Hermann Minkowski (1864 – 1909)

Space and Time *Minkowski's Papers on Relativity* 



#### Willem de Sitter (1872 – 1934)

Leiden Observatory Archives, Willem de Sitter papers (WdS)



"It is known that geometry assumes, as things given, both the notion of space and the first principles of constructions in space.

• • •

This leads us into the domain of another science, of physic [sic], into which the object of this work does not allow us to go to-day."

– G. F. Bernhard Riemann (1854)

G. F. B. Riemann, "On the Hypotheses which lie at the Bases of Geometry," *Göttingen Inaugural Lecture* (1854).

 $\frac{\mathrm{d}t}{\mathrm{d}\tau}$ 

# Time measurement is only *locally* valid; time is a strictly-local phenomenon.

– Albert Einstein (1905)

A. Einstein, "On the Electrodynamics of Moving Bodies," Annalen der Physik 17, 891 (1905).

$$=\frac{1}{\sqrt{1-\frac{v^2}{c^2}}}$$

$$\mathrm{d}\tau^2 = -c^2 \mathrm{d}t$$


# Space and time are locally-orthogonal coordinates in *the absolute* [4D] *world*.

– Hermann Minkowski (1908)

ict

H. Minkowski, "Space and Time," A Lecture delivered in Cologne (21 Sept. 1908).

 $t^2 + \mathrm{d}x^2 + \mathrm{d}y^2 + \mathrm{d}z^2$ 




#### An exact solution to the field equations

$$ds^{2} = -dr^{2} - R^{2} \sin^{2}\left(\frac{r}{R}\right) \left[d\psi^{2} + \sin^{2}\left(\psi\right)d\theta^{2}\right] + c^{2}dt^{2}$$

### The Universe is finite, yet has no boundary.

A. Einstein, "Cosmological Considerations in the General Theory of Relativity," SPAW, 142 (1917).





### Einstein's cosmological considerations


The objective of this paper is not simply to present an historical overview of Einstein's cosmological considerations, but to discuss the central role they played in shaping the paradigm of relativistic cosmology. This, we'll show, was a result of both his actions and, perhaps more importantly, his inactions. Accordingly, discussion won't simply be restricted to Einstein's considerations, as we'll analyse relevant contributions to the relativistic expansion paradigm during the approximately twenty years following Slipher's first redshift measurements in 1912. Our aim is to shed some light on why we think some of the things we do, with the idea that a better understanding of the reasoning that fundamentally influenced the common idea of our expanding universe might help to resolve some of the significant problems that modern cosmology now faces; and we eventually use this knowledge to probe the foundations of the standard model. Much of the information we present, including many of the historical details, we expect will be news to modern practitioners.

Cosmology is in a crisis state unprecedented in the history of science. For while we now possess a model describing the large-scale evolution of our Universe whose parameters have been constrained with significant precision [unsurprisingly not actually so in hindsight], many aspects of the universe that this model seems to describe simply don't meet our theoretical expectations—and a number of 'big questions' have therefore arisen along with the great scientific progress that's taken place this past century.

D. Janzen, "Einstein's cosmological considerations" (2014); arXiv:1402.3212 [physics.hist-ph].

Daryl Janzen, Lecturer in Physics, Dept. of Physics and Engineering Physics, Univ. of Saskatchewan







EINSTEIN only assumes *three*-dimensional space to be finite. It is in consequence of this assumption that in (2A)  $g_{44}$  remains 1, instead of becoming zero with the other  $g_{\mu\nu}$ . This has suggested the idea 1) to extend EINSTEIN'S hypothesis to the four-dimensional time-space. We then find a set of  $g_{\mu\nu}$  which at infinity degenerate to the values [(2B)]. Moreover we find the remarkable result, that now no "world-matter" is required.

– Willem de Sitter (31 March 1917)

W. de Sitter, "On the relativity of inertia. Remarks concerning Einstein's latest hypothesis," *KNAW Proceedings* **19**(2), 1217 (1917).

1) The idea to make the four-dimensional world spherical in order to avoid the necessity of assigning boundary-conditions, was suggested several months ago by Prof. EHRENFEST, in a conversation with the writer. It was, however, at that time, not further developed.

$$\left|\begin{array}{cccccccccc}
0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 \\
\end{array}\right|$$
(2B)



#### Willem de Sitter had this very important critical insight...

"We thus find that in [Einstein's metric] the time has a separate position. That this must be so, is evident a priori. ... Such a fundamental difference between the time and the space-coordinates seems to be somewhat contradictory to the complete symmetry of the field-equations..."

– Willem de Sitter (31 March 1917)

W. de Sitter, "On the relativity of inertia. Remarks concerning Einstein's latest hypothesis," *KNAW Proceedings* **19**(2), 1217 (1917).



#### but then he got confused and found no physical interpretation.

"All this shows that the postulate of the invariance of  $g_{\mu\nu}$  at infinity [emphasis added] has no real physical meaning. It is purely mathematical."

– Willem de Sitter (31 March 1917)

W. de Sitter, "On the relativity of inertia. Remarks concerning Einstein's latest hypothesis," *KNAW Proceedings* **19**(2), 1217 (1917).

W. de Sitter, "Einstein's theory of gravitation and its astronomical consequences. Third paper," *MNRAS* 78, 3 (1917).

Also see the 1918 *KNAW* paper covering a 30 June 1917 lecture on the next slide (quoted reference).

Einstein

de Sitter

If we neglect all pressures and other internal forces, and if we  
suppose all matter to be at rest, then the tensor 
$$T_{\mu\nu}$$
 becomes  
(5)  $T_{44} = g_{44}\rho$ , all other  $T_{\mu\nu} = 0$ ,  
 $\rho$  being the density in natural measure. We can put  
(6)  $\rho = \rho_0 + \rho_1$ ,  
where  $\rho_0$  is the average density of the world-matter. If  $\rho_0$  i  
positive, then  $\rho_1$  may be positive or negative; but in the latter cas  
the numerical value must not exceed  $\rho_0$ .  
Nov. 1917. Gravitation, and its Astronomical Consequences. 7  
If we wish to neglect gravitation, we must neglect  $\rho_1$ , and  
take  $\rho_0$  constant. The equations (3) then become \*  
(7)  $\begin{cases} G_{ij} - (\lambda + \frac{1}{2}\kappa\rho_0)g_{ij} = 0.\\ G_{44} - (\lambda + \frac{1}{2}\kappa\rho_0)g_{44} = -\kappa\rho_0g_{44}.\end{cases}$   
These can be satisfied by the  $g_{\mu\nu}$  implied by the line-element  
(8A)  $ds^2 = -dr^2 - R^2 \sin^2 \frac{r}{R} [d\psi^2 + \sin^2\psi d\theta^2] + c^2 dt^2$ ,  
if  
(9A)  $\kappa\rho_0 = 2\lambda$ ,  $\lambda = \frac{1}{R^2}$ .  
This is Einstein's new solution.  
The equations are also satisfied by  
(8B)  $ds^2 = -dr^2 - R^2 \sin^2 \frac{r}{R} [d\psi^2 + \sin^2\psi d\theta^2] + \cos^2 \frac{r}{R} c^2 dt^2$ ,  
if  
(9B)  $\rho_0 = 0$ ,  $\lambda = \frac{3}{R^2}$ ;





## Another exact solution to the field equations

$$ds^{2} = -dr^{2} - R^{2} \sin^{2}\left(\frac{r}{R}\right) \left[d\psi^{2} + \sin^{2}(\psi)d\theta^{2}\right] + \frac{\cos^{2}\left(\frac{r}{R}\right)c^{2}dt^{2}}{R}$$

### A Universe with '*zero average* density' (i.e., net zero energy) seems absurd—or is it?

W. de Sitter, "On the curvature of space," KNAW Proceedings 20(1), 229 (1918).

- "... we have  $\rho_0 = 0$  and consequently all  $T_{\mu\nu} = 0$ ."
  - $\varrho_0 \equiv$  "average matter density"
  - Willem de Sitter (30 June 1917)






### Distance-induced time dilation [f(r)] in the de Sitter-metric

$$ds^{2} = -dr^{2} - R^{2} \sin^{2}\left(\frac{r}{R}\right) \left[dy^{2}\right]$$
$$ds^{2} \equiv d\tau^{2} = \cos^{2}\left(\frac{r}{R}\right)c^{2}dt^{2}$$

$$\frac{\mathrm{d}t}{\mathrm{d}\tau} = \frac{1}{\cos r}$$


 $\psi^{2} + \sin^{2}(\psi) d\theta^{2} + \cos^{2}\left(\frac{r}{R}\right)c^{2} dt^{2}$ 


$$\underbrace{(R=c=1)}_{\text{normalized}}$$



animation...

The distinct physical interpretations of Einstein's and de Sitter's " $g_{44}$ "\*






\* (by modern convention, now " $g_{00}$ ")

The distinct physical interpretations of Einstein's and de Sitter's " $g_{44}$ "\* Note that this model derives independently of the EFE from symmetry (i.e., the 3-sphere) and local orthogonality of space and time (Minkowski).

asynchronous time  $[\tau(r)]$ 



animation completed

View the animation (YouTube) https://youtu.be/kswjXqnNEs0




Change quality to 1080p.

 $t \equiv \tau(0)$ 

$$+\cos^{2}\left(\frac{r}{R}\right)c^{2}dt^{2}$$
  
de Sitter

\* (by modern convention, now " $g_{00}$ ")



### Time reversal and energy polarity


From any observer's arbitrary location in the Cosmos, local proper time in the fundamentally-invisible antipodal volumetric half of the Universe is systemically (i.e., azimuthally averaged) *relativistically reversed* as compared to local proper time in the adjacent visible half, which is bounded by the cosmological redshift horizon (see next slide).

The average matter density of both halves (A, B) is equal, so the two energy *magnitudes* are equal:  $|E_A| = |E_B|$ . However, *relativistically*,  $E_A = -E_B$ ; therefore the net cosmic energy is zero:  $E_A + E_B = 0$ .

A remarkably-insightful speculation dating back fifty years was on the right track! K. Mullis, "Cosmological Significance of Time Reversal," Nature 218, 663 (1968).

An idea arising from mathematical physics, which involves symmetry and the Planck equation (E = hv), is that time reversal yields negative energy.


photo credit: Kent Clemenco



Kary Mullis

Cosmic antipodal energy polarity is a temporal *relativity* phenomenon.

Same galaxy, different observer location. Radial arrows represent local proper time. Locally, the laws of physics (e.g., the second law of thermodynamics) are identical.



Relativistically, the net energy of the Universe is always zero: (E - E = 0)from the point of view of an observer at any arbitrary Cosmic location.






### From Chapter 19, "The Future of Time"


Lee Smolin

"To make further progress in cosmology (and in fundamental physics as well), we need a new conception of a law of nature, valid on the cosmological scale, which avoids the fallacies, dilemmas, and paradoxes and answers the questions that the old framework cannot address." - Lee Smolin, Time Reborn (2013), p. 240.

"The nature of space and time is *the* basic subject in physics." Online Lecture: The Nature of Space and Time (0:01:15 / 1:28:25)









### **Redshift-distance relationship**

R is the normalized cosmic radius; r is the distance from the observer;  $\chi$  is the "cosmic latitude"  $[0 \le \chi \le \pi]$ .\*



\*  $\chi \equiv 0$  (where  $dt \equiv d\tau_0$ ) is defined for an observer at any arbitrary location in the Universe (e.g., the Milky Way).



• • •

$$\ln[1]:= \theta = \frac{2.0}{3600} * \frac{\pi}{180}; (* 2 \text{ arcseconds in radi})$$

$$\ln[2]:= \rho = \frac{15}{\theta}; (* \text{ ESTIMATE (15 kly) per (2 \text{ arcsec})})$$

$$estimated intrinsic diameter and expression \left[\rho * \left(1 - \frac{1}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(0.00(1)\right)^{-\frac{1}{2}}\right)^{-\frac{1}{2}} * \left(0.00(1)\right)^{-\frac{1}{2}} * \left(1 - \frac{1}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(0.00(1)\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(0.00(1)\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(0.00(1)\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.08 + 1)^2}\right)^{-\frac{1}{2}} * \left(1 - \frac{\pi}{(0.0$$

### lans \*)

conds @ z = 0.08 \*) galaxy nuclei empirically-observed apparent diameter \* 1\*^-6, 2] (\* estimated Cosmic radius in Gly \*) as per new theta-z formula

t horizon in Gly \*)

e Universe in ly<sup>3</sup> \*)

in ly<sup>3</sup> for 10<sup>12</sup> galaxies \*)

0<sup>12</sup> galaxies in ly \*)

117

٦٦

Ð

# The Cosmic Radius

In the previous slide, the theta-z predictive formula was used to determine the Cosmic radius (R)as the single unknown. In that formula, the two directly-measurable values are the redshift (z) and the apparent angular diameter ( $\theta$ ) of an observed distant object. The remaining empirical variable, which must be estimated, is the intrinsic diameter ( $\delta$ ) of that object, and  $\delta/\theta = \rho$ , where this Euclidean effective radius (*rho*) is illustrated on slide 137.

According to the empirical data in the form of visually-analyzed SDSS galaxy images, ubiquitous galaxy nuclei are all of approximately the same intrinsic diameter, which diameter is independent of the widely-varying (by about an order of magnitude) host-galaxy size. Based on observation of the Milky Way and M31 (Andromeda), that diameter is estimated to be about 15 kly.

As concerns *absolute* measurements, the single free empirical parameter shared by all predictive formulas is the estimated Cosmic radius R. As such, its accurate measurement by the above method warrants a precision astronometric program (e.g., using the light curves of galactic nuclei) and a more precise measurement of  $\delta$  than the round estimate of 15 kly.

Though not shown for notebook brevity,  $\rho(0.08) \approx 1.55$  Gly. At this redshift, the range in Petrosian radius is  $7 \le r_P \le 17$ , which correlates to a range in *typical* galaxy diameter of  $\rho \cdot 2r_P \sim 25 - 250$  kly. This is a "reality check" (with good results) on the estimated value of  $\rho$ , from which we derive R.









### Prior "redshift-independent" distance measurements

The astronomical reference literature dating back more than a half-century includes numerous "redshift-independent" extragalactic distance measurements, including error bars. For nearby galaxies (out to an estimated 100 Mly) the primary measurement method is the period-luminosity relation of Cepheid variables. For more distant galaxies, in which Cepheids cannot be resolved, the assumed characteristics of Type IA supernovae, the Tully-Fisher relation (an adopted correlation between galaxy luminosity and rotational velocity), and the Faber-Jackson relation (an adopted correlation between elliptical-galaxy luminosity and nuclear stellar velocity dispersion) have been the primary such measurement methods.

In reality, all of these extragalactic distance indicators were tied to the purported 'Hubble constant,' which was interpreted as a rigorous empirical constraint on the redshift-distance relationship, recently thought to have 1% error bars.<sup>1</sup> Extragalactic distance measurements contradicting such a generally-accepted (albeit false) 'empirical law' were thus considered to be 'impossible.'

As it is consistent with SDSS and other statistically-significant empirical data, the new redshiftdistance relationship is expected to yield absolute distance measurements that are correct to the accuracy of the estimated Cosmic radius R. Corrections to prior distance measurements can be attributed to the 'impossibility' of those measurements to have disagreed with the 'Hubble law.' 1. C. L. Bennett, D. Larson, J. L. Weiland, & G. Hinshaw, "The 1% Concordance Hubble Constant," ApJ 794, 135 (2014).







$$\underbrace{\overset{r}{6.5 \text{ Mly}} = 4.1 \text{ Gly} \cdot \cos^{-1}\left(\frac{1}{z+1}\right)}_{r}$$

•	🐲 Hubb
In[1]:=	R = 4.1*^9; (* estimated Cosmic ra
In[2]:=	r = 6.5*^6; (* 2 Mpc is about 6.5
In[3]:=	Clear[z]; (* solve for redshift (
In[4]:=	$\{z\} = z /.$ Solve $\left[r = R * ArcCos \left[\frac{1}{z+1}\right]\right]$
Out[4]=	$\left\{1.25669 \times 10^{-6}\right\}$
In[5]:=	2.998*^5 * z (* redshift expressed
Out[5]=	<b>0.376757</b> (Yields $y \sim 0$ flat green line in sli


Given  $R \sim 4.1$  Gly, then in the 1929 Hubble diagram, which extends to  $r = 2^+$  Mpc:


redshift-distance relationship

```
ble Diagram cz.nb
adius is 4.1 Gly *)
                                                         D
Mly *)
                                                         2
(z), given R and r *)
                                                         Z
                                                        Д.
- , z
                                                        Ξ
d as a recession velocity cz in km s<sup>-1</sup> *)
lide-56 Hubble diagram, given its "VELOCITY" scale.)
                                                        P
                               125% ▶
```

# Mapping a finite, boundaryless 2-space...

Every point in a two-dimensional homogeneous, finite, boundaryless space (S<sup>2</sup>) can be mapped by an infinite set of great circles ( $S^1$ ). A more intuitive map of the space is the union of two disks ( $S^2$ ) whose boundary points are duplicate coordinates, whereby a midline (e.g.,  $\overline{AB}$  where A and B map and identical point) represents one such great circle. Ignoring terrestrial asymmetry, line segments  $\overline{0A}$  and  $\overline{0B}$  are linear projections of semicircles having 'identical' length to similar paths on the Equator.






Line  $\overline{AB}$  represents a circle.



# Mapping a finite, boundaryless 3-space


Every point in a three-dimensional homogeneous, finite, boundaryless space (S<sup>3</sup>) can be mapped by an infinite set of spheres (S<sup>2</sup>). A more intuitive map of the space is the union of two balls (S<sup>3</sup>) whose exterior boundary points are duplicate coordinates. The interior midline  $\overline{AB}$  represents a cosmic great circle as A and B map the identical point. The set of interior points uniquely map all of cosmic space, less the set of duplicated points on the map's boundary. Line segments  $\overline{0A}$  and  $\overline{0B}$  are linear projections of semicircles having identical length to similar paths on the boundary.



Line  $\overline{AB}$  represents a circle.



# A complete cosmological map of the S<sup>2</sup> surface *locally* defined by Galactic latitude b = 0 (i.e., the Galactic plane)



Graphic by Fabio Basile after an original rendering by Hollin Calloway (2005).


 $z = \infty$ @ horizon  $(\chi = \pi/2)$ 





 $\cos \chi$ 

Distance



### Comparison of redshift-distance relationship

124

(redshift) Ζ


## Reference to the local Universe

 $-122 \pm 8 \text{ km} \cdot \text{s}^{-1}$ , the Doppler shift equivalent of the former being v/c = -0.001.

According to the following Aitoff projection sky map, such measured galaxy blueshifts, caused by relative-motion ("peculiar-velocity") Doppler shifts overshadowing distance redshift, are ubiquitous within the footprint of redshift surveys that faithfully record them. The bar charts on the next slide (127) indicate that measured heliocentric blueshifts rarely exceed  $|z| \sim 0.001$ , which puts constraints on Doppler-induced variation between measured galaxy shift (±) and *distance* redshift.

The two redshift-magnitude graphs (128–129), with corresponding distance scales in light years, use M31 as a reference standard candle. Its apparent visual magnitude of 3.4 is in accord with a nominal r-band value of 2.5 mags. Low-redshift data ( $z \le 0.12$ ), which includes aperture-limited measurements for the nearest galaxies, is sourced from SIMBAD with other data from SDSS. The two graphs compare the Euclidean ACDM model to the new, Riemannian 'de Sitter' model. The redshift value of M31 in both graphs is based on its  $\sim 2.6$  Mly redshift-independent distance.

Our closest neighbor, M31 (Andromeda Galaxy) has a mean redshift-independent distance of 0.784 Mpc (~2.6 Mly). According to the consensus ACDM cosmological model, with parameter values ( $H_0$ ,  $\Omega_M$ ,  $\Omega_\Lambda$ ) shown repeatedly in prior graphs, that distance measurement corresponds to a distance redshift z(d) = 0.000182. Its blueshifted (–) heliocentric relative velocity is recorded as  $-300 \pm 4 \,\mathrm{km} \cdot \mathrm{s}^{-1}$ , while its Galactocentric relative velocity (reflecting the Sun's orbital motion) is










## ~6500 *blueshifted* galaxies...

126




\* Arecibo General Catalog  $\ddagger \sim 5.5 \text{k}$  galaxies (2.5% of  $\sim 221 \text{k}$  w/quality  $\geq$  3)





\* Actual measured *blueshift* is  $-10^{-3}$ .



# PART IV – NEW PREDICTIVE EQUATIONS

Click to go back to Table of Contents...

The tool implementing the mediation between theory and practice, between thought and observation, is mathematics. Mathematics builds the connecting bridges and is constantly enhancing their capabilities. Therefore it happens that our entire contemporary culture, in so far as it rests on intellectual penetration and utilization of nature, finds its foundations in mathematics.


For us there is no ignorance, especially not, in my opinion, for the natural sciences. Instead of this silly ignorance, on the contrary let our fate be: "We must know, we will know."

– David Hilbert, Preeminent 20th-century mathematician (1862–1943)



. . .

Source: *Translation of an address given by David Hilbert in Königsberg*, Fall 1930; translation by Amelia and Joe Ball.



### **David Hilbert**



## The topological unit 3-sphere ( $S^3$ ) has a finite, boundaryless volume.

line element:  $ds^2 = d\psi^2 + \sin^2\psi$ Denotes the topological 3-sphere; not an exponent!  $S^3 = \int_0^{\pi} \mathrm{d}\psi \int_0^{\pi} \sin\psi$  $S^3 = 4\pi \int_0^\pi \sin^2 \psi$ volume:  $S^3 = 2\pi(\psi - \cos\psi)$  $\psi \to \chi$  $S^{3} = 2\pi (\chi - \cos \chi \sin \chi)$  $S^3(\chi)$ :


$$\psi \left( \mathrm{d}\theta^2 + \sin^2\theta \, \mathrm{d}\phi^2 \right)$$

$$\mathbf{v} \, \mathrm{d}\boldsymbol{\theta} \, \int_0^{2\pi} \sin \boldsymbol{\psi} \, \sin \boldsymbol{\theta} \, \mathrm{d}\boldsymbol{\phi}$$

$$d\psi$$

$$\psi\sin\psi\big)\Big]_0^\pi=2\pi^2$$


"continued on next slide"




### Redshift-volume function...

$$S^{3}(\chi) = 2\pi (\chi - \zeta)$$
$$\begin{cases} \cos \chi = \frac{1}{(z+1)} \\ \sin \chi = \sqrt{1 - \cos^{2}} \\ S^{3}(z) = 2\pi \begin{cases} \cos^{-1} \chi \\ \cos^{-$$

 $\cos\chi\sin\chi)$ 






### Volume-element function...


$$S^{3}(z) = 2\pi \left[ \cos^{-1} \left( \frac{1}{z+1} \right) - \left( \frac{1}{(z+1)^{2}} - \frac{1}{(z+1)^{4}} \right)^{\frac{1}{2}} \right]$$
$$u = (z+1)^{-1} \qquad \frac{\mathrm{d}u}{\mathrm{d}z} = -(z+1)^{-2} = -u^{2}$$

$$S^{3}(z) = 2\pi \left[ \cos^{-1} u - \left(u^{2} - u^{4}\right)^{\frac{1}{2}} \right]$$
$$\frac{dS^{3}}{dz} = 2\pi \left[ \frac{-1}{\sqrt{1 - u^{2}}} \frac{du}{dz} - \frac{1}{2\sqrt{u^{2} - u^{4}}} \left( 2u \frac{du}{dz} - 4u^{3} \frac{du}{dz} \right) \right]$$



### Volume-element function...

$$\frac{dS^{3}}{dz} = 2\pi \left[ \frac{-1}{\sqrt{1-u^{2}}} \frac{du}{dz} - \frac{1}{2\sqrt{u^{2}-u^{4}}} \left( 2u \right) \right]$$
$$= 2\pi \left[ \frac{u^{2}}{\sqrt{1-u^{2}}} - \frac{1}{2\sqrt{u^{2}-u^{4}}} \left( -2u^{3} + \frac{1}{2\sqrt{u^{2}-u^{4}}} \left( -2u^{3} + \frac{1}{\sqrt{1-u^{2}}} + \frac{1}{\sqrt{u^{2}-u^{4}}} \left( u^{3} - 2u^{5} + \frac{1}{\sqrt{1-u^{2}}} \left( u^{2} - u^{4} \right) \right] \right]$$
$$= 4\pi \left[ \frac{1}{\sqrt{1-u^{2}}} \left( u^{2} - u^{4} \right) \right]$$






$$\frac{dS^{3}}{dz} = 4\pi \left[ \frac{1}{\sqrt{1 - u^{2}}} \left( u^{2} - u^{4} \right) \right] \qquad \left\{ u = (z + 1)^{-1} \right\}$$
$$\frac{dS^{3}}{dz} = \frac{4\pi}{\sqrt{1 - (z + 1)^{-2}}} \left( \frac{1}{(z + 1)^{2}} - \frac{1}{(z + 1)^{4}} \right) \quad \underline{R^{3} \text{ units}}$$


$$S^{3}(z) = C_{V} \cdot 2\pi \left[ \cos^{-1} \left( \frac{1}{z+1} \right) - \left( \frac{dS^{3}}{dz} = C_{dV} \cdot \frac{4\pi}{\sqrt{1-(z+1)^{-2}}} \left( \frac{1}{(z+1)^{-2}} \right) \right]$$

Volume-element function...

$$\left\{u=\left(z+1\right)^{-1}\right\}$$




Here,  $C_V$  and  $C_{dV}$  are data-dependent arbitrary scaling constants.



### Theta-z relationship (apparent size of a standard rod)


Jump back to The Cosmic Radius.



Here,  $C_{\delta}$  is an arbitrary *scaling* constant that is proportional to the intrinsic diameter of the standard rod (a class of astrophysical object understood to be of approximately the same intrinsic size) under consideration.



$$\rho = \mathcal{R} \sin \chi = \left(1 - \frac{1}{(z+1)^2}\right)^{\frac{1}{2}}$$
$$\theta(z) \approx \frac{1}{\rho}$$
$$\theta(z) = C_{\delta} \left(1 - \frac{1}{(z+1)^2}\right)^{-\frac{1}{2}} \text{ radians}$$





$$S^{3} = 4\pi \int_{0}^{\pi} \sin^{2} \psi \, d\psi = 4\pi$$
$$S^{2} = \frac{d}{d\chi} S^{3} = 4\pi \sin^{2} \chi$$

$$S^{2} = 4\pi \left( 1 - \frac{1}{(z+1)^{2}} \right) \frac{R^{2}}{(z+1)^{2}}$$

Redshift-magnitude function...

 $\pi \int_0^{\pi} \sin^2 \chi \, \mathrm{d}\chi \qquad (\psi \to \chi)$ 






$$F(S^{2}) = \frac{L}{S^{2}} = \frac{L}{4\pi \left(1 - \frac{1}{(z+1)^{2}}\right)}$$

$$F(z) = \frac{L}{4\pi \left(1 - \frac{1}{(z+1)^2}\right) \cdot (z+1)^4} = \frac{L}{4\pi \left[(z+1)^4 - (z+1)^2\right]}$$

Redshift-magnitude function...

# Time dilation factor : $(z+1)^{-2}$ Curvature factor : $(z+1)^{-2}$




$$F(z) = \frac{L}{4\pi \left[ \left( z + 1 \right)^4 - \left( z + 1 \right)^4 \right]}$$


$$m = C_M - 2.5 \cdot \log(b)$$

$$m(z) = C_M - 2.5 \cdot \log \left( \frac{1}{4\pi \left[ (z+1)^4 - (z+1)^2 \right]} \right)$$
 mags

Here, C<sub>M</sub> is an arbitrary scaling constant that is proportional to the absolute magnitude of the standard candle under consideration.

Redshift-magnitude function...






# Redshift-magnitude function

If one wishes to incorporate IGM extinction (light dimming due to absorption and scattering by the intergalactic medium), which is *approximated*<sup>\*</sup> to be a linear function of radial distance, then

$$m(z) = C_M - 2.5 \cdot \log \left( \frac{1}{4\pi \left[ \left( z+1 \right)^4 - \left( z+1 \right)^2 \right]} \right) + \varepsilon_\lambda \cos^{-1} \left( \frac{1}{z+1} \right) \text{ mags}$$
  
IGM extinction

Here,  $\varepsilon_{\lambda}$  is a constant, which may vary with observed wavelength ( $\lambda$ ); it is correlated to the average density of the IGM over the line of sight to the target object. This was set to *zero* for the graphed functions.

\* Such approximation does not take into account that IGM extinction is a function of photon wavelength, which varies over the light path due to persistent redshift.



## PART V – GALAXY SPACE DENSITY

Click to go back to Table of Contents...

# Space-density of active galactic nuclei

To counteract any survey selection effects, data for ~108k AGN is sourced from the NASA/IPAC Extragalactic Database (NED); SDSS data constitutes about 72% of the graphed NED AGN dataset, with the balance sourced from numerous other surveys.


A small percentage of galaxies host an active galactic nucleus (AGN), which has a much higher luminosity than normal; the brightest AGN are observed at very high redshift (z > 6), so these galaxies are an ideal observable with which to confront theoretical predictions of galaxy space density, including testable assumptions concerning galaxy evolution over lookback time. According to the data and the new cosmological model, AGN constitute a fixed portion of the local regional galaxy population (on large scale) throughout the Universe. According to the ACDM model, AGN population density has increased by over four orders of magnitude since the presumed 'primordial universe.'

# NED AGN


# Click *image* for source data, here for precompiled datafile.

You h		1		•	
Activity 7	Type: AGN		1 1 1		
NED	results for your specified	Inarame	ters		
	courts for your specific	r par anno			
	0		1 20 2		
08214	objects found in NED.	· · · ·		1 A 1	
	/	1		2 Garage	
bjects 1	-100 displayed. Page 1 of 1083	3 ≫ Go to p	age: go		
		-			
	a the second	1			
	CE LIST	1 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4			
SOUR	Object list is sorted				
		on RA and Object Type	Dec Redshift	Row No.	
SOUR	Object list is sorted	<b>O</b> bject		No. 1	
Row No.	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627	Object Type QSO QSO	Redshift 1.356000 1.143400	No. 1 2	
Row No.	<b>Object list is sorted</b> <b>Object Name</b> FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226	Object Type QSO QSO QSO	Redshift 1.356000 1.143400 1.331200	No. 1 2 3	
Row No.	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627	Object Type QSO QSO	Redshift 1.356000 1.143400	No. 1 2	
<b>Row</b> No.	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226 HELLAS2XMM 26900016	Object Type QSO QSO QSO G	Redshift 1.356000 1.143400 1.331200 1.314000	No. 1 2 3 4	
<b>Row</b> <b>No</b> .	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226 HELLAS2XMM 26900016	Object Type QSO QSO QSO G G	Redshift 1.356000 1.143400 1.331200 1.314000	No. 1 2 3 4	
<b>Row</b> <b>No</b> .	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226 HELLAS2XMM 26900016 2MASSi J0000029-350332	Object Type QSO QSO QSO G G	Redshift 1.356000 1.143400 1.331200 1.314000	No. 1 2 3 4	
<b>Row</b> <b>No.</b> 1 2 3 4 5	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226 HELLAS2XMM 26900016 2MASSi J0000029-350332	Object Type QSO QSO QSO G G G	Redshift 1.356000 1.143400 1.331200 1.314000 0.508000 1.248000 1.776658	No. 1 2 3 4 5 108210 108211	
<b>Row</b> <b>No.</b> 1 2 3 4 5	Object list is sorted Object Name FBQS J0000-0202 2QZ J000001.3-303627 2QZ J000001.7-312226 HELLAS2XMM 26900016 2MASSi J0000029-350332 2QZ J235958.3-283223 SDSS J235958.66-011225.2 SDSS J235958.72+003345.3	Object Type QSO QSO QSO G G	Redshift 1.356000 1.143400 1.331200 1.314000 0.508000 1.248000	No. 1 2 3 4 5	

P








z (redshift)

### Cumulative AGN population ( $\sum AGN \leq z$ ) vs. volume models



# Space-density of Seyfert galaxies

Any uniquely-identifiable subset of the galaxy population that is expected to constitute some fixed percentage of the total galaxy population can be used to confront the volume function with empirical data. One such example are galaxies classified as Seyfert galaxies, which are spiral galaxies with unusually bright nuclei that produce broad emission lines. Referencing NED and using a similar procedure as that used to analyze the general AGN population, the redshift population distribution of Seyfert galaxies is graphed subsequently...


### Click *image* for source data, here for precompiled datafile.

	Type: Sy				1.00
ED	mogulto for your monifi				2
ED		od honom	otores		
	results for your specifi	eu param	eters:		
			· · ·	8.62	
585	objects found in NED.				
	2 have been a second	· 🥖 ·	1		1.00
bjects	1-100 displayed. Page 1 of 1	46 <u>&gt;&gt;</u> Go to p	page: go		•
		i inc			
SOUD					
SOUR	CELIST				1
	CE LIST				
		d on RA and	1 Dec		
	CE LIST Object list is sorte	d on RA and	d Dec		
	Object list is sorte				
		Object	d Dec Redshift	Row	
	Object list is sorte			Row No.	
No. 1	Object list is sorte Object Name LQAC 000-025 001	Object Type G	<b>Redshift</b> 1.314000	No. 1	
<b>No.</b> 1	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508	Object Type G G	Redshift 1.314000 0.102441	No. 1 2	
No. 1	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508 2MASS J00001172+0523175	Object Type G G G	Redshift 1.314000 0.102441 0.040000	No. 1	
<b>No.</b> 1	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508	Object Type G G	Redshift 1.314000 0.102441	No. 1 2	
Row No. 1 2 3 4 5	<b>Object list is sorte</b> <b>Object Name</b> LQAC 000-025 001 2MASS J00001161+0101508 2MASS J00001172+0523175 LQAC 000-025 002 LQAC 000-025 003	Object Type G G G G G	Redshift 1.314000 0.102441 0.040000 0.995000	No. 1 2 3 4	
No. 1 2 3 4 5	Object list is sorte           Object Name           LQAC 000-025 001           2MASS J00001161+0101508           2MASS J00001172+0523175           LQAC 000-025 002           LQAC 000-025 003	Object Type G G G G	Redshift 1.314000 0.102441 0.040000 0.995000 1.610000	No. 1 2 3 4	
No. 1 2 3 4 5	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508 2MASS J00001172+0523175 LQAC 000-025 002 LQAC 000-025 003	Object Type G G G G G	Redshift 1.314000 0.102441 0.040000 0.995000 1.610000 0.556000	No. 1 2 3 4 5	
No. 1 2 3 4 5 14581 14582	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508 2MASS J00001172+0523175 LQAC 000-025 002 LQAC 000-025 003	Object Type G G G G G	Redshift 1.314000 0.102441 0.040000 0.995000 1.610000 0.556000 0.738000	No. 1 2 3 4 5 14581 14582	
<b>No.</b> 1	Object list is sorte Object Name LQAC 000-025 001 2MASS J00001161+0101508 2MASS J00001172+0523175 LQAC 000-025 002 LQAC 000-025 003	Object Type G G G G G	Redshift 1.314000 0.102441 0.040000 0.995000 1.610000 0.556000	No. 1 2 3 4 5	


P

14,585 Seyferts


**NED Seyferts** 











# PART VI – CREDITS AND AFTERWORD

Note: Addenda 1–2 (Parts VII & VIII) follow.

Click to go back to Table of Contents...

# SDSS-I/II Acknowledgment (Legacy data)

Funding for the SDSS and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, the Max Planck Society, and the Higher Education Funding Council for England. The SDSS Web Site is www.sdss.org.

The SDSS is managed by the Astrophysical Research Consortium for the Participating Institutions. The Participating Institutions are the American Museum of Natural History, Astrophysical Institute Potsdam, University of Basel, University of Cambridge, Case Western Reserve University, University of Chicago, Drexel University, Fermilab, the Institute for Advanced Study, the Japan Participation Group, Johns Hopkins University, the Joint Institute for Nuclear Astrophysics, the Kavli Institute for Particle Astrophysics and Cosmology, the Korean Scientist Group, the Chinese Academy of Sciences (LAMOST), Los Alamos National Laboratory, the Max-Planck-Institute for Astronomy (MPIA), the Max-Planck-Institute for Astrophysics (MPA), New Mexico State University, Ohio State University, University of Pittsburgh, University of Portsmouth, Princeton University, the United States Naval Observatory, and the University of Washington.



# SDSS-III Acknowledgment (BOSS data)

Funding for SDSS-III has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, and the U.S. Department of Energy Office of Science. The SDSS-III web site is www.sdss3.org.

SDSS-III is managed by the Astrophysical Research Consortium for the Participating Institutions of the SDSS-III Collaboration including the University of Arizona, the Brazilian Participation Group, Brookhaven National Laboratory, Carnegie Mellon University, University of Florida, the French Participation Group, the German Participation Group, Harvard University, the Instituto de Astrofisica de Canarias, the Michigan State/Notre Dame/JINA Participation Group, Johns Hopkins University, Lawrence Berkeley National Laboratory, Max Planck Institute for Astrophysics, Max Planck Institute for Extraterrestrial Physics, New Mexico State University, New York University, Ohio State University, Pennsylvania State University, University of Portsmouth, Princeton University, the Spanish Participation Group, University of Tokyo, University of Utah, Vanderbilt University, University of Virginia, University of Washington, and Yale University.

### SCIENCE BLOG FROM THE SDSS

News from the Sloan Digital Sky Surveys

http://blog.sdss3.org

NASA's Astrophysics Data System;

the VizieR catalogue access tool, CDS, Strasbourg, France Reference: *A&AS* 143, 23;

the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.



VIN340









### This research has made use of

# JPL HORIZONS System;

### the SIMBAD database, CDS, Strasbourg, France Reference: *A&AS* 143, 9;

### 

### 

"The age of 10<sup>18</sup> years for rich compact clusters of galaxies may be shortened somewhat by considering certain interactions between galaxies that lead to more inelastic and resonant encounters between galaxies. Unless, however, far greater efficiency for the transfer of energy and momentum is postulated for such interactions than is compatible with our present-day knowledge of physical phenomena, the age of rich spherically symmetrical and compact clusters of galaxies is clearly greater than 1015 years."1


– Fritz Zwicky, Preeminent 20th-century astrophysicist (1898–1974)

"If ever a competition were held for the most unrecognized genius of twentieth century astronomy, the winner surely would be Fritz Zwicky."

- Steven Soter & Neil deGrasse Tyson, Cosmic Horizons: Astronomy at the Cutting Edge. New York: The New Press, 2000.

1. F. Zwicky, "The Age of Large Globular Clusters of Galaxies," Pub. Astron. Soc. Pacif. 72 (428), 365 (1960).

### This is a reasonable scientific statement (c. 1960) based on empirical evidence:




Fritz Zwicky



# A DEFINITIVELY-FALSIFIED MODEL

Ø



This isn't just wrong—it's "spherically wrong." (i.e., the same from every perspective, as per Fritz Zwicky)

Dark Energy Dark Matter **Ordinary Matter** 

26.4% 4.4% 69.2%

Plank Collaboration, *Plank* 2015 Results. XIII. Cosmological Parameters; arXiv:1502.01589 [astro-ph.CO]

### The greater meaning of this document beyond its scientific content


I want to argue that clear thinking, combined with a respect for evidence — especially inconvenient and unwanted evidence, evidence that challenges our preconceptions — are of the utmost importance to the survival of the human race\* in the twenty-first century, and especially so in any polity that professes to be a democracy.

– Alan Sokal, Professor of physics at New York University Professor of mathematics at University College London

\* One must concede that there is such a thing as "dangerously stupid."



Source: A. Sokal, "What is science and why should we care? — Part I," *Scientia Salon* (26 March 2014).



Alan Sokal





Tycho Brahe' (1546 - 1601)

 $^{2}\mathcal{O}$ 

# James Edward Gunn, "Father" of the SDSS

"In 1987 Gunn proposed putting an array of CCDs on a 2.5m-telescope and using it for both images and spectra, scanning the entire visible sky in about five years and building an enormous data archive which could be used for far more than his main interest, determining the three-dimensional structure of the universe of galaxies. This ultimately became the Sloan Digital Sky Survey, and Gunn devoted a large portion of his career to building it and making it work."


Source: http://www.phys-astro.sonoma.edu/brucemedalists/Gunn/index.html

### The perfect candidate for the NOBEL PRIZE IN PHYSICS for *real* contributions to science.

J. E. GUNN ET AL., "THE 2.5 m TELESCOPE OF THE SLOAN DIGITAL SKY SURVEY," THE ASTRONOMICAL JOURNAL, 131:2332–2359, 2006 April

\* Tycho Brahe's accurate astronomical data allowed Johannes Kepler to discover that the planets moved in elliptical orbits, which led to Isaac Newton's universal law of gravitation. James Gunn's work proves to be of a similar nature.





157

Photo 2009 by Brian Wilson • brianwilsonphotographer.com •

Impromptu, hand-held, amateur video footage taken during the talk and Q&A:

https://youtu.be/KIQmVdgMv14

I should not have used the phrase, "catastrophic failure of the model" during my talk, as this may be interpreted as a subjective opinion, and it could engender strong negative emotions in certain individuals. People should come to that conclusion on their own from the evidence.

I thank Dr. César Augusto Zen Vasconcellos for the opportunity to present this research at IWARA 2018.





I thank the entire SDSS Team over many years (from DR4 in 2005 to present). I am privileged to have had the opportunity to benefit from their life's work and they share directly in the achievement of scientific discovery that arises from it.

In addition, I thank the engineers, computer scientists, executive management and support staff of Apple Computer, Wolfram (Mathematica), Google, Wikimedia, Adobe, Microsoft (SQL Server), Oracle (MySQL), and PremiumSoft (Navicat) for providing the tools and services that performed essential roles in this research.


Although prior analysis of CMBR and supernovae data, of which I am aptly critical, may have been flawed, it is clear that the related projects have contributed in no small measure to this research. I thank the teams of engineers, technicians, and astronomers, in particular the legion of NASA/JPL staff and NASA contractors, who made those challenging observational enterprises possible.

This research is not the work of one person; it is the success of many thousands.









James Webb Space Telescope Earliest Launch: March 2021

- 6.5-m 18-segment primary mirror
- Near Infrared Camera (NIRCam)  $0.6 \le \lambda \le 5$  microns
- Near Infrared Spectrograph (NIRSpec)  $0.6 \le \lambda \le 5$  microns
- Mid-Infrared Instrument (MIRI)  $5 \le \lambda \le 28$  microns
- Fine Guidance Sensor/Near Infrared Imager and Slitless Spectrograph (FGS/NIRISS)  $0.8 \le \lambda \le 5$  microns


There is no such thing as the "early Universe"; JWST will see that cosmological creation is an eternal process and that the Universe is similar yet different everywhere, as is true for Earth.












# PART VII – ADDENDUM 1: RELATED COSMOLOGICAL ISSUES

Click to go back to Table of Contents...

eternal dynamic equilibrium, having a static volume ( $\dot{R} = 0$ ):

- 1. Given the mutual gravitational attraction of the total cosmic matter content over an arbitrary amount of time, what prevents general gravitational collapse?
- 2. Given that a primordial hot, dense state of the Universe is a false inference based on the fallacious idea of an expanding universe, then 'Big Bang nucleosynthesis' of the light-element nucleotides (1H, 2H, 4He, 3He, 7Li, 7Be) is certainly another fallacy. As deuterium (2H) is destroyed in the interiors of stars faster than it is produced, its observed cosmic abundance (i.e., its source) requires explanation.
- 3. Given that galaxy redshifts are accurately modeled by relativistic time dilation induced by uniform, isotropic cosmic spacetime curvature, their prior canonical interpretation as universal expansion is falsified; thus, no primordial compressed, hot, dense state of the Universe approximately 14 Gyr ago can be presumed. Accordingly, a *ubiquitous and continuous energy* conversion process is required to explain the observed cosmic microwave background.
- 4. What is the nature of purported 'dark matter'?

- Three cornerstones of the Big-Bang theory and 'dark matter'
- Four important issues require resolution in the context of a 'de Sitter' Universe in a state of









### Dynamical cosmic mass-energy redistribution

amount of time, what prevents general gravitational collapse?

The geometric model of relativistic time applicable to the cosmological Riemannian 3-sphere (S<sup>3</sup>) is *similarly* applicable in the case of spacetime geometry encountered in the strong-field limit (i.e., a black hole). It is apparent that the present canonical conceptualization of such phenomenon as a compact spatially-localized *object* having a 'singularity' at coordinate r = 0 is unphysical. Such a "hole" *necessarily* connects two antipodal regions of cosmological spacetime having relativistically-reversed time coordinates. It is a quasi-stable configuration of spacetime-geometry, which *cosmological structure* is sustained by dynamic mass-energy *flow* through that structure. Every locally-mass-energy-*absorbing* black hole has a corresponding *cosmologically-antipodal* locally-mass-energy-*emitting* "white hole." Such white holes are empirically observed as regions of mass-energy emission far exceeding that which is possible through thermonuclear fusion. While mass-energy conservation holds cosmologically, it does not hold locally in the vicinity of such holes, which transport mass-energy between cosmological antipodes.

**1.** Given the mutual gravitational attraction of the total cosmic matter content over an arbitrary





The reciprocal cosmological and energetic relationships between the terminals of a wormhole.

Image courtesy of NRAO/AUI/NSF


# Wormholes

It proves to be the case that the foundations of general relativity are inconsistent with the Newtonian conception of a gravitational '*equipotential* surface,' which surface is simply defined by a finite coordinate radius about an idealized point source: Having a strictly-locally-defined geometric definition according to these foundations, each among the set of distinct local proper time coordinates  $\tau_p$ , associated with the neighborhood of each point p on such surface, excepting antipodal points, cannot have the identical geometric definition (i.e., they are not parallel in  $R^4$ ). This distinction correlates to a symmetric time dilation between the  $\tau_p$  with associated measurable (already measured yet unrecognized) empirical phenomena in the weak field.\*

This empirically-verifiable incremental improvement to canonical general relativity radically alters the interpretation of the strong-field limit and eliminates the dubious notion that a physical 'spacetime singularity,' where the known laws of physics break down, exists at coordinate r = 0. Rather, as has been long suspected, in particular by the pure mathematics community, spacetime must be smooth and continuous everywhere; catastrophic gravitational collapse leads to the creation of a physically-real so-called "wormhole" connecting distinct regions of spacetime as illustrated in the preceding slide.

\* Details to be discussed in a pending series of journal letters by A. F. Mayer.





# Cosmic jets

### 2. Where do observable conditions exist for the production of deuterium (<sup>2</sup>H)?

A very high-temperature plasma (~ $10^9$  K or ~0.1 Mev) of free protons and neutrons that cools rapidly is understood to be a process that leads to the nucleosynthesis of the light elements. The necessary high temperatures for the required reaction sequences to take place are not found in stars and deuterium production requires an environment of high energy coupled with low density. The observed existence of the light elements cannot be due to normal processes in stellar evolution, particularly because typical stellar evolution involves destruction of deuterium.

Observed jets, as exemplified in the foregoing image of Radio Quasar 3C175, constitute an ideal environment for nucleosynthesis of the light elements. It is understood that such jets emanate from "white holes," which may be characterized as cosmological-scale particle accelerators. Such jets are the likely progenitors of new (i.e., spiral) galaxies, so one would expect to find an unlikely abundance of <sup>2</sup>H at the core of spiral galaxies (e.g., the Milky Way). This is exactly the case, although its presence was attributed to an ad hoc phenomenon.<sup>1</sup>

1. D. A. Lubowich et al., "Deuterium in the Galactic Centre as a result of recent infall of low-metalicity gas," *Nature* **405**, 1025 (2000).

**3.** What is the source of the cosmic microwave background (CMB)?

As it is now conclusively determined that the Universe is not expanding, it is certain that the CMB does not have a 'primordial origin.' Accordingly, one must identify some continuous and contemporary source for these photons. Controversy about the origins of the CMB is not new.<sup>1</sup> It should now be clear that published analyses of data strongly confirming the Big-Bang theory were fabricated; however, atypical analyses of the same data that has been largely ignored to date has yielded important insights:

The cosmic microwave background is often called the echo of the Big Bang, but recent research suggests that some of its features might have their origins much closer to home. Although most cosmologists think that the tiny variations in the temperature of the background are related to quantum fluctuations in the early universe, Glenn Starkman and colleagues at CERN and Case Western Reserve University in the US have now found evidence that some of these variations might have their roots in processes occurring in the solar system. If correct, the new work would require major revisions to the standard model of cosmology. ... "Each of these correlations could just be an accident," says Starkman. "But we are piling up accident on accident. Maybe it is not an accident and, in fact, there is some new physics going on."<sup>2</sup>


1. H.-J. Fahr and M. Sokaliwska, "Remaining Problems in Interpretation of the Cosmic Microwave Background," *Phys. Res. Int.* **2015**, 503106.

2. E. Cartlidge, "Doubts cast over map of the cosmos," *Physics World* **18(1)**, 5 (2005).





### WMAP full-sky temperature maps (linear scale from -200 to $+200 \mu$ K) – WMAP Science Team



"The Internal Linear Combination Map is a weighted linear combination of the five WMAP frequency maps. The weights are computed using criteria which minimize the Galactic foreground contribution to the sky signal. The resultant map provides a low-contamination image of the CMB anisotropy." – NASA LAMBDA









### Some claimed interpretations of WMAP data (source verbatim)

### The WMAP science team has...

- 4. ...nailed down the curvature of space to within 0.4% of "flat" Euclidean.
- universe.
- of atoms) is 24.0%

Source: https://map.gsfc.nasa.gov

An important paper with a prophetic title that challenged these interpretations:

U. Sawangwit & T. Shanks, "Is everything we know about the universe wrong?" Astronomy & Geophysics 51, 5.14 (2010).

3. ...determined the universe to be 13.77 billion years old to within a half percent.

5. ...determined that ordinary atoms (also called baryons) make up only 4.6% of the

6. ...completed a census of the universe and finds that dark matter (matter not made up

7. ...determined that dark energy, in the form of a cosmological constant, makes up 71.4% of the universe, causing the expansion rate of the universe to speed up. -"Lingering doubts about the existence of dark energy and the composition of the universe dissolved when the WMAP satellite took the most detailed picture ever of the

cosmic microwave background (CMB)." - <u>Science Magazine 2003, "Breakthrough of the Year" article</u>







### "Excess microwave emission"

We now know that there is no such thing as a 'primordial Big-Bang photon,' so it is evident that the much-touted analyses of COBE-, WMAP-, and Plank-satellite data, purporting to 'prove' various aspects of the Big-Bang theory were fabrications, not unlike the systemic fabrication of data uncovered during the 2008 'financial crisis.' The problem in physics is one of "group-think," arguably driven by the agendas of a few key 'authorities,' who have received high accolades.

From the following quotation, it is clear that the galactic foreground emission was actually never understood, thus requiring unrestrained ad hoc speculation. Then the process of sanitizing the "contaminated" (subjectively bad) empirical data to produce 'science data' (subjectively good) involved *arbitrarily* removing theoretically-problematic objective empirical data.


> The cause of observed inner galaxy excess microwave emission is assumed to be synchrotron emission from highly relativistic electron-positron pairs produced by [ad hoc] dark matter particle annihilation as more conventional sources have been ruled out.<sup>1, 2</sup>


1. D. P. Finkbeiner, "WMAP Microwave Emission Interpreted as Dark Matter Annihilation in the Inner Galaxy," (2 Sep. 2004); arXiv:astro-ph/0409027.

 $\rightarrow$  Phys. Rev. D 76, 083012 (2007)

2. D. P. Finkbeiner, "Microwave Interstellar Medium Emission Observed by the Wilkinson Microwave Anisotropy Probe," *Astrophys J.* **614**, 186–193 (2004).









Also, in 2003 Hans Kristian Eriksen of the University of Oslo and his co-workers presented more results that hinted at alignments. They divided the sky into all possible pairs of hemispheres and looked at the relative intensity of the fluctuations on the opposite halves of the sky. What they found contradicted the standard inflationary cosmology—the hemispheres often had very different amounts of power. But what was most surprising was that the pair of hemispheres that were the most different were the ones lying above and below the Ecliptic, the plane of the earth's orbit around the sun. This result was the first sign that the CMB fluctuations, which were supposed to be cosmological in origin, with some contamination by emission in our own galaxy, have a solar system signal in them—that is, a type of observational artifact.<sup>1</sup>

— astrophysicists have found that the plane of the solar system threads itself through hot and cold spots in the cosmic microwave background, suggesting that some of the variations in the latter are not caused by events that took place in the early universe.<sup>2</sup>

1. G. D. Starkman and D. J. Schwarz, "Is the Universe Out of Tune?" *SciAm* (Aug. 2005), p. 52. 2. E. Cartlidge, "Doubts cast over map of the cosmos," *Physics World* **18(1)**, 5 (2005).

### Vitally-important conventionally-disregarded empirical evidence from WMAP data



The large-angle (low- $\ell$ ) correlations of the cosmic microwave background exhibit several statistically significant anomalies compared to the standard inflationary cosmology.

• • •

• • •

We have shown that the planes defined by the octopole are nearly aligned with the plane of the Dopplersubtracted quadrupole, that three of these planes are orthogonal to the ecliptic plane, with normals aligned with the dipole (or the equinoxes), while the fourth plane is perpendicular to the supergalactic plane. Each of these correlations is unlikely at 99% C.L., and at least two of them are statistically independent. We have also seen that the ecliptic threads between a hot and a cold spot of the combined Doppler-subtracted-quadrupole and octopole maps—following a node line across about 1/3 of the sky, and separating the three strong extrema from the three weak extrema of the map.

We find it hard to believe that these correlations are just statistical fluctuations around standard inflationary cosmology's prediction of statistically isotropic Gaussian random  $a_{\ell m}$ 's. That the quadrupole-octopole correlation just happened to increase by ~5 when the quadrupole was Doppler-corrected seems particularly unlikely. The correlation of the normals with the ecliptic poles suggest an unknown source or sink of CMB radiation or an unrecognized systematic. If it is a physical source or sink in the inner solar system it would cause an annual modulation in the time-ordered data.


<u>Physical correlation of the CMB with the equinoxes is difficult to imagine, since the WMAP satellite has no knowledge of the inclination of the Earth's spin axis.</u> (all underlined emphasis added)

D. J. Schwarz, G. D. Starkman, D. Huterer & C. J. Copi, "Is the Low-ℓ Microwave Background Cosmic?" *Phys. Rev. Lett.* **93**, 221301 (2004); arXiv:astro-ph/0403353.

# The primary cause of secular astrophysical spin-down (i.e., "rotation braking")

The principle of "relativistic temporal geometry," which has been shown to apply cosmologically, must similarly apply to the local gravitational field. The mathematical expression of the latter phenomenon is off-topic herein. However, to summarize it, just as a redshift occurs due to the homogeneous and isotropic curvature of cosmological space between widely-separated regions at the same Newtonian potential, the same principle applies over the curvature incurred in any orbit in a gravitational field, including such orbit at constant radius around an ideal point-mass. Due to a (geometric) distinction in proper time at each point on the orbit, by analogy, one may consider a gravitational field to have a kind of 'resistance,' causing subtle energy loss for orbiting bodies in the weak field that manifests in the form of electromagnetic radiation, which radiation can be thought of as a kind of 'heat' due to that 'resistance.' The stronger the field, and the greater the velocity of motion through it, the greater the energy-dissipation effect, per unit mass.

Constituent mass-elements of a rotating astrophysical body effectively 'orbit' in the gravitational field of the total source mass, with the highest velocity occuring at its equator. It follows that the temperature of the radiation produced by the effective 'gravitational resistance' incurred by that rotational motion is greatest at the equator and falls off to zero at the poles. Note that energy conservation correlated with this phenomenon requires the secular spin-down of all rotating astrophysical bodies. This *universal* 'gravitational-resistance' spin-down is particularly noticeable for pulsars, in which case that effect is conventionally attributed to magnetic dipole radiation.



### Predicted observable phenomena

The foregoing assertion concerning the fundamental instability of gravitational systems and the associated radiative phenomenon is subject to empirical verification: The Sun represents 99.86% of Solar System mass, it has an equatorial sidereal rotation period of about 600 hours (longer at higher latitude), and its equatorial plane is inclined 7.25 degrees to the Ecliptic. Due to that obliquity, the South Pole Telescope points toward the solar equatorial plane for six months per year, and then away from it. One may expect to see a correlated annual modulation in microwave temperature in that data (see next slide, *left* diagram).


For a satellite co-orbiting with the Earth at L2 in the Ecliptic (e.g., WMAP), the obliquity of the Earth's spin axis to the Ecliptic (23.44°) causes a similar observable effect correlated to the equinoxes (see next slide, *right* diagram). From astronomical records dating back several millennia, the long-term increase in the mean length-of-day has been established to be about 1.8 ms per century,<sup>1</sup> This secular spin-down correlates to an average continuous energy dissipation of 2.8 terawatts. The conventional explanation for terrestrial rotational braking is that Earth's rotational energy is being dissipated as "tidal friction" and also "tidal acceleration" of the Moon. The Sun, rather than Earth, is the Moon's dominant gravitational field and it's differential 'resistive' quality on the Earth-Moon system causes observed lunar acceleration.

1. F. R. Stephenson, L. V. Morrison2, & C. Y. Hohenkerk, "Measurement of the Earth's rotation: 720 BC to AD 2015," *Proc. R. Soc. A* **472**, 20160404.










SDSS terapixel image of the night sky. Limited to the "Northern Galactic Cap," this image represents just a portion of the WMAP "full-sky map," below.



**Internal Linear Combination** 

### Every point in this image is a galaxy.




Image credit: M. Blanton and the SDSS-III

*Every galaxy* emits microwave radiation, as does our Galaxy (red foreground, below).

> This is just  $\sim 10^{6}$ galaxies of  $\sim 10^{11}$ cosmologically.

Northern Galactic Cap









With  $\sim 10^{11}$  stars orbiting its center, the Galaxy is an obvious source of the identical radiation correlated to the secular spin-down of the Sun and Earth, also each star's orbital motion in the galactic gravitational field contributes to the radiation. Hence, the huge Galactic signal that had to be removed from the popularized WMAP "Internal Linear Combination Map," including inexplicable "excess microwave emission."

Given a total population on the order of 10<sup>11</sup>–10<sup>12</sup> galaxies, the Cosmos presents the observer with about  $10^4$ – $10^5$  galaxies per micro-steradian, that representing  $\sim 10^{-7}$  of the celestial sphere. With each galaxy locally producing the same microwave radiation as observed for the Milky Way (e.g., the K-Band Map at 23 GHz), it is clear that this presents the observer with a nearlyisotropic microwave background, with slightly higher temperature for regions of higher density.

The foregoing implies that the phenomenon of "gravitational radiation" is ubiquitous, observably manifesting in the electromagnetic spectrum. Verification of the predicted annual modulation in time-ordered  $\mu$ -wave data would show that the COBE, WMAP, and Plank satellites were de facto 'gravitational wave' detectors and that LIGO and similar instruments are useless white elephants. The purported measurement of "gravitational waves" according to their heretofore canonical theoretical description, and leading to the 2017 Nobel Prize in Physics, was at best bad science, and at worst a fraud, as has been argued by various authors (see next slide)...

Ubiquity of 'gravitational radiation'







### Selected supporting references jeopardizing the celebrated LIGO 'discovery'

D. Christopoulos, "My deepest disappointment for Nobel Prize Physics 2017," RG (Dec 2017).

black holes observation," RG (Mar 2016).

B. R. De, "Casebook of Bibhas De—The Mystery of the Hanford Empaths," eBook (7 Jan 2018).

B. R. De, "Unchallenged privilege: The billion-dollar trilateral gravitational-wave discovery scam," 2nd Ed., Bibhas De (Aug 2017).

RG (Oct 2017).

J. Horgan, "Is the Gravitational-Wave Claim True? And Was It Worth the Cost?," blogs.scientificamerican.com (12 Feb 2016).

Despite claims of authenticity, a telling description of how an artificial signal, indistinguishable from 'science data,' was a distinct possibility:

M. Mahin, "LIGO Doubts Will Persist Unless Replication Occurs," futureandcosmos.blogspot.com (19 Feb 2016).

X. Mei, Z. Huang, P Ulianov, & P. Yu, "LIGO Experiments Cannot Detect Gravitational Waves by Using Laser Michelson Interferometers," J. Mod. Phys. 7, 1749 (2016).

S. Sims, "Problems with the LIGO gravitational wave discovery," Plasma Pics (15 Mar 2016).

- J. Creswell, S. von Hausegger, A. D. Jackson, H. Liu, & P. Naselsky, "On the time lags of the LIGO signals," (9 Aug 2017); arXiv:1706.04191v2. J. Creswell, S. von Hausegger, A. D. Jackson, H. Liu, & P. Naselsky, "Comments on our paper, 'On the time lags of the LIGO signals'," (27 Jun 2017).
- D. Christopoulos, "A detailed critical review of reported event GW150914 that LIGO/VIRGO collaboration announced as gravitational waves and
- W. W. Engelhardt, The LIGO-VIRGO Miracle: A contemplation on the detection of gravitational waves in three different places on August 14, 2017,"
- J. Kanner & A. Weinstein, "The Astrophysicists Who Faked It—The inside story of the gravitational wave signal injection," Naultilus (3 Nov 2016).





# 'Empirical evidence' can be illusory

### **4.** What is the nature of 'dark matter'?

Purported empirical evidence for 'dark matter' includes apparent excess (otherwise-undetectable) gravitational mass required to bind galaxy clusters, to produce observed gravitational lensing, and to account for spiral-galaxy rotation curves. Rather than the common perception that such rotation curves are "flat," they actually exhibit radial acceleration (increasing orbital velocity) as per observation of the hydrogen 21-cm line (i.e., orbiting gas) extending far beyond the optical disk.


Observed radial acceleration of spiral galaxy rotation curves is readily explained in the context of the gravitational tidal forces on the disk produced by the host cluster; details are available at gravitysim.net, in particular, this 15-minute video (immediate viewing recommended). A definitive solution to the perplexing winding problem concerning spiral galaxies is also put forward.

Prior calculations based on observed gravitational lenses suffered from application of a faulty redshift-distance relationship and Cosmic geometry (Euclidean versus Riemannian), causing misinterpretation of empirical observables (next slide). In the subsequent slide, redshift distances for the Einstein ring and background galaxy, which are based the new model herein, indicate no non-luminous excess mass within the boundary of the ring.









### Field of view: $0.13 \times 0.13$ arcminutes $R_{\rm eff} = 2.16 \pm 0.13$ " $z_{FG} = 0.3223$ $z_{BG} = 0.5812$

A. S. Bolton, et al., "The Sloan Lens Acs Survey. I. A Large Spectroscopically Selected Sample Of Massive Early-Type Lens Galaxies," ApJ 638, 703 (2006).

#### Einstein Ring Gravitational Lens SDSS J073728.45+321618.5

Credit: NASA, ESA, A. Bolton (Harvard-Smithsonian CfA) and the SLACS Team

• • • 🌻 Einstein Ring.nb  $\ln[1] = \theta_{\rm E} = \frac{2.16^3}{2+3600} * \frac{\pi}{180} ; (* \text{ radians } *)$  $\ln[2]:= G = 6.67408^{6*} - 11; (* m^{3} kg^{-1} s^{-2} *)$  $\ln[3] = c = 299792458^9; (* m s^{-1} *)$  $\ln[4] = R = 4.1^{2*^9} * 9.46073^{15}; (* m *)$  $\ln[5]:= D_{L} = R * \operatorname{ArcCos}\left[\frac{1}{0.3233 + 1}\right];$  $\ln[6]:= D_{s} = R * \operatorname{ArcCos}\left[\frac{1}{0.5812 + 1}\right];$  $ln[7]:= D_{LS} = D_{S} - D_{L};$ In[8]:= Clear [M]  $\ln[9]:= \{M\} = M / . \text{ Solve} \left[ \theta_E = \sqrt{\frac{4 G \star M}{c^2}} \frac{D_{LS}}{D_L \star D_C}, M \right];$ In[10]:= M = UnitConvert[Quantity[M, "Kilograms"], "SolarMass"] Out[10]=  $6.62313 \times 10^{11} M_{\odot}$ 

The here-calculated value of  $6.6 \times 10^{11} \text{ M}_{\odot}$  inside the Einstein ring is consistent with conventional matter; no mysterious, non-luminous 'dark matter' need be attributed to the mass of this galaxy or similar ones.

Z.	$\Lambda CDM^* (D_A)$	'de Sitter'	*
0.3223	3.1753 Gly	2.9278 Gly	$H_0 = 69.6$
0.5812	4.4810 Gly	3.6331 Gly	$\Omega_{\rm M} = 0.28$ $\Omega_{\Lambda} = 0.71$
$D_{LS}$	0.7934 <b>Gly</b>	0.7053 Gly	$\Box c_{\Lambda} = 0.7 T$





### Explanation of following Mathematica notebook and correlated graph

The Coma cluster contains  $\sim 1k$  galaxies observed within apparent diameter  $\sim 100'$ . For R = 4.1 Gly,  $1^{rad}$  corresponds to ~872 Mly at the reference redshift for Coma ( $z_{ref} = 0.0234$ ); so the Coma radius (50') correlates to  $r_c = 12.7$  Mly. Relative to  $z_{ref}$ , this radius gives near/far boundaries in redshift space for the Coma cluster of  $z \approx 0.0227/0.0241$ . Accordingly, the highlighted red box represents a ~25 Mly-diameter ball roughly centered on NGC 4874.

Assuming that most of the galaxies in the graph are "clustered" inside that ball, then one must interpret most of their redshifts as radial velocities  $\sim c(z_n - z_{ref})$ , shown by the scale on the right. If one furthermore assumes that such unlikely configuration is dynamically stable, one must then invent invisible gravitating matter having no electromagnetic signature whatsoever, and having a mass within the ball many times that of the visible baryonic matter. Some time ago, this purported 'dark matter' was given the prophetic British acronym FAIRIE DUST: "Fabricated Ad hoc Invention Repeatedly Invoked in Efforts to Defend Untenable Scientific Theory."

Alternatively, one may interpret the measured redshifts as being predominantly *cosmological*; then the plotted "Coma-cluster" galaxies are distributed along a filament  $\sim 25$  Mly in diameter with a ~360 Mly extent. That filament has a higher-to-lower center-to-periphery density gradient that is highlighted by the horizontal (radial-distance scale) background-shading gradient in the graph. Observing Coma, one is looking down a filament, rather than at a roughly-spherical distribution.








• • •

In[1]:= R = 4100`; (\* estimated Cosmic radius in Mly \*) In[2]:= z<sub>ref</sub> = 0.0234`; (\* Coma-cluster redshift as per SIMBAD \*)  $\ln[3] = r_c = 3000$ ; (\* observed cluster radius of 50 arcmins in arcsecs \*)  $\ln[4]:= \rho = R \left( 1 - \frac{1}{(z_{ref} + 1)^2} \right)^{\frac{1}{2}} (* 1 \text{ radian at } z_{ref} \text{ in Mly } *)$ Out[4]= 871.741  $\rho = \frac{\pi}{180 + 3600}$  1\*^6 (\* ly per arcsecond @  $z_{ref}$  for top x-axis \*) Out[5]= 4226.32  $\frac{\pi}{180 + 3600} \rho \ (* \ r_c \ converted \ to \ radians \rightarrow Mly \ *)$ Out[6]= 12.679  $\ln[7] = D_m = R * \operatorname{ArcCos} \left[ \frac{1}{0.0234 \pm 1} \right] (* z = 0.0234; \operatorname{Coma-cluster} z_{ref} \text{ distance in Mly } *)$ Out[7]= 878.446  $\ln[8]:= Solve \left[ D_m - r_c = R * ArcCos \left[ \frac{1}{\pi + 1} \right], z \right] (* matching Coma-cluster far redshift *)$ Out[8]=  $\{ \{ z \rightarrow 0.0227167 \} \}$  $\ln[9]:= \text{Solve}\left[D_m + r_c == R * \operatorname{ArcCos}\left[\frac{1}{z+1}\right], z\right] (* \text{ matching Coma-cluster far redshift } *)$ Out[9]=  $\{ \{ z \rightarrow 0.024094 \} \}$ 

```
🏘 Coma.nb
```

125% ▶



# NGC 4889

Adam Block/Mount Lemmon SkyCenter/University of Arizona

HST detail (next slide).

### Coma cluster center

star

# NGC 4874


 $1200 \text{ arcsecs} = 1/5 \times 100 \text{ arcmins}$ 

~5 Mly @ z = 0.0234



NASA, ESA, and the Hubble Heritage Team (STScI/AURA)

# Coma cluster detail



100-arcmin variable diameter to scale

roughly spherical clump; then "galaxy cluster" conveys the tacit presumption that the observed group of galaxies is similarly structured—to be sure, one does not refer to a long queue of persons as a *cluster*. By inspection and analysis, the name "Coma cluster" is a misnomer; one should rather call that region of sky the "Coma filament." The observed galaxies are distributed over a long and narrow region of space as illustrated toscale at left. To simply assume that they are "clustered" within the region of the red ball, which requires very high relative velocities that could only be induced by gravitating 'dark matter,' is both arbitrary and unphysical. Being based on the incorrect expanding-universe model,

all previous galaxy-evolution models are equally wrong. Galaxies and their groups did not form by local accretion (i.e., "clustering") of expanding matter.

By convention, the word "cluster" generally refers to a







# PART VIII – ADDENDUM 2: RELEVANT HISTORY, ETC.

Click to go back to Table of Contents...

# Brilliant Blunders

#### 

#### FROM DARWIN TO EINSTEIN

Colossal Mistakes by Great Scientists That Changed Our Understanding of Life and the Universe

#### Μ Α

"Physicists sometimes tend to ignore the history of their subject. After all, who cares who discovered what, as long as the discoveries are made widely known? In a few cases, however, discoveries are of such magnitude, that understanding the path that had led to such insights, including the correct attribution, can be of great value."1

About the book: "Breakthroughs require the willingness to embrace risks and to accept errors as potential portals of discovery."

2 1. Mario Livio, "Lost in translation: Mystery of the missing text solved," Nature 479, 171 (2011); free HubbleSite version.

- Astrophysicist and author, Mario Livio



Mario Livio





In 1921, a young Belgian mathematics postdoc and seminarian by the name of Georges Henri Lemaître wrote an essay entitled *God's First Three Declarations*.\* The author stated that this 1921 essay was "an attempt to describe scientifically the first verses of Genesis." This biblically-inspired essay, discovered in the archives of the Catholic University of Leuven in the late 20<sup>th</sup> century, is the actual root of the Big-Bang theory. Lemaître (July 1894 – June 1966) was ordained as a Catholic priest in 1923.

\* Footnotes: "Les trois premières paroles de Dieu." The manuscript is reproduced in Stoffel (1996), pp. 107–111. Lemaître's religious views are discussed in Lambert (1997).

Source: Helge Kragh, *Matter and Spirit in the Universe* (Imperial College Press, London, 2004), p. 141.

Click the book cover for Journal for the History of Astronomy review article by E. McMullin (2005).



#### Georges Lemaître

HISTORY OF MODERN PHYSICAL SCIENCES - VOL. 3

Matter

Scientific and Religious Preludes to Modern Cosmology

Helge Kragh Aarhus University, Denmark



### In June 1925, Lemaître met with Hubble at Mt. Wilson.<sup>1,2</sup>



1. John Farrell, *The Day Without Yesterday: Lemaître, Einstein, and the Birth of Modern Cosmology*, New York: Thunder's Mouth Press, 2005, p. 78. 2. Jeremiah P. Ostriker & Simon Mitton, *Heart of Darkness: Unraveling the Mysteries of the Invisible Universe*, Princeton: Princeton Univ. Press, 2013, p. 68.



- (1) references Mt. Wilson(2) references Caltech
- (2) references Caltech



### The Lemaître-Hubble connection

In 1926, the year following his special trip to Mt. Wilson, Georges Lemaître submitted a paper to *Annales de la Société scientifique de Bruxelles* that was published in 1927 in Belgium (in French), so it had a limited audience: "A Homogeneous Universe of Constant Mass and Increasing Radius accounting for the Radial Velocity of Extra-galactic Nebulæ."<sup>1</sup>

Lemaître's visit to Mt. Wilson was motivated by his nascent cosmological theory and, three years after this initial meeting in California, Edwin Hubble and Lemaître both attended the 1928 IAU (International Astronomical Union) conference in Holland, giving Hubble an additional opportunity to meet with Lemaître.<sup>2</sup> It is unlikely that Lemaître did not discuss the details of his fantastical expanding-universe theory with Hubble when they met on both occasions (1925 and 1928).

Referencing a 1926 *ApJ* paper by Hubble,<sup>3</sup> Lemaître's paper provided a cosmic expansionvelocity estimate of 625 km s<sup>-1</sup> Mpc<sup>-1</sup>, whose inverse implied a 'Hubble time' (i.e., the age of the modeled expanding universe) of about 1.6 Gyr, this at a point in history when geologists were rapidly coming to the conclusion that the planet Earth had existed for at least 2–3 Gyr.

M. l'Abbé G. Lemaître, "Un Univers Homogéne de Masse Constante et de Rayon Croissant...," Annales de la Société Scientifique de Bruxelles A47, 49 (1927).
 Sidney van den Bergh, "Discovery of the Expansion of the Universe," J. Roy. Astron. Soc. Can. 105(5), 197 (2011); arXiv:1108.0709 [physics.hist-ph].
 Edwin Hubble, "Extra-Galactic Nebulae," ApJ 64, 321 (1926).



#### Grace Lillian Burke Hubble

Edwin Hubble's wife, Grace, was a wealthy and influential socialite,\* having inherited two large fortunes, the first from her father and the second from her first husband. That elite status was considerably amplified, as it was concurrent with the Great Depression, and it likely had a strong influence on her husband's media-driven reputation.

\* "Grace Hubble, a brilliant woman with a keen wit, led an extraordinary life at the center of Caltech's intellectual community, with members of the British émigré community in Los Angeles and among Hollywood's elite."

Solution Content of Content of Antice And Antice Ant



#### "...a man whose life is cloaked in pathological lies..."

[The screenplay] "Hubble" is the magnificent story of one of history's greatest and most flawed geniuses and the even more magnificent universe he sought to map. In 1931, Edwin Hubble became the most famous man in the world. He was heralded as the greatest astronomer since Galileo. His discoveries had an irrevocable impact on both Einstein's Theory of Relativity and religious interpretations of the origins of heaven and earth. But Hubble was a haunted man, dogged by mysterious secrets from the past and by enemies that threatened to destroy everything. How could a man who spoke with a British accent, wore a cape, and carried a cane be from Missouri? Why did none of his stories of his past match the claims of others? How could his wife Grace knowingly perpetuate all of this? Driven by intense ambition and a longing for something that was lost long ago, a man whose life is cloaked in pathological lies paradoxically discovers one of science's greatest and most enduring truths [that turns out to be one of science's greatest and most enduring cons and a plagiarism of a young priest's very-misguided, religiously-inspired fantasy.]

Tribeca Film Institute (Alfred P. Sloan Partnership), Comment on the screenplay, "Hubble," by S. L. Otto (2005).



Quotation from reference 3 of 4 in E. Hubble's 1929 PNAS paper; the italics are in the original:



Curtist seems to have been the first to realise the cosmical importance of the occurrence of Novæ in spiral nebulæ. By assuming equality in absolute magnitude of galactic and spiral Novæ he concluded that the latter, being apparently ten magnitudes fainter, are about one hundred times as far away as are the former. Estimating the mean distance of galactic Novæ to be 100,000 light years, he reached the conclusion that the spirals are galactic systems in size comparable to our own, and that the closest spirals are millions of light-years away.

<sup>2</sup> *Journal of the Washington Acad. of Sc.*, **9**, 217 (1919) & L.O.B., **300**, 108 (1917) (Lick Observatory Bulletin)

An excerpt from the referenced L.O.B. article appears on the following slide...

Oddly, Edwin Hubble is credited for original work by other people...

#### <sup>3</sup> Mon. Not. R. Astr. Soc., 85, 1925 (865–894).



#### Oddly, Edwin Hubble is credited for original work by other people...

The occurrence of these novae in spiral nebulae must be regraded as a fact bearing very directly on the theories of the constitution of the spirals. It seems to me that they furnish weighty evidence in favor of the well known "island universe" theory of the spiral nebulae. ...

... There is thus an average difference of ten magnitudes between galactic novae and spiral novae. Now all the evidence available assigns a great distance to the galactic novae. If we assume equality of absolute magnitude for galactic and spiral novae, then the latter, being apparently ten magnitudes the fainter, are of the order of one hundred times as far away as the former. That is, the spirals containing the novae are far outside our stellar system [i.e., the Milky Way]; and these particular spirals are undoubtedly, judging from their relatively greater angular diameters, the nearer spirals. ...

Herber D. Curtis U.S. Shipping Board Navigation School San Diego, California, September 8, 1917.

Issued October 16, 1917.

Quotation: Lick Observatory bulletin 300 (1917), p. 110.


SERVATORY BULLE NUMBER 300

#### HREE NOVAE IN SPIRAL NEBULAE

1915 June 3, H. C. O. 1 inch Cooke lans; limiting magnitude 12.5; an object is suspected.

VOLUME IX & AVA W AN AN AN TO JON 13 4 10-1, 4, 1939

1905 Feb. 11, H. C. O. Drappet telescope: limiting magnitude for the center of the note.
1907 Feb. 16, Yorken Bisneb Brane; the nebula is 5° from the center of the plate; on the net of norm.
1915 Jan. 13, H. C. O. Linch Corke less; limiting magnitude 155 Mar. 20, L. O. Grossley. The plate was taken of N. G. 4587 and he nova are aboven near the edge of the plate. Own the near of the distance from the center of the plate in the neav are aboven near the edge of the plate. Own the neaver are aboven near the edge of the plate. Own the near are aboven near the edge of the plate. Own the neaver are aboven near the edge of the plate. Own the near are aboven near the edge of the plate. Own the near are aboven near the edge of the plate. Own the superared in the outer part of this spiral at some time prior to March 2, 1914, 24" east and 110". South of the nucleus. Thoogh Nova B is recorded on only the one plate, there is no doubt that it is a home fade sellar image; it shows the same petuliarities of factors are at the time is fully furnished me by Directors Frest, inde 14.
1915 Jare 3, H. C. O. Linch Corke less; limiting magnitude 15.
1915 Jare 3, H. C. O. Linch Corke less; limiting magnitude 15.



Click for document.

#### Oddly, Edwin Hubble is credited for original work by other people.

this was published five years earlier by Carl Wilhelm Wirtz in Germany: De Sitters Kosmologie und die Radialbewegungen der Spiralnebel. Von C. Wirtz. zunehmender Entfernung ganz erheblich anwächst." Doppler shift, cz] of the spiral nebulae grows very considerably with distance." Carl W. Wirtz, "De Sitter cosmology and the radial motion of the spiral nebulae," *Astronomische Nachrichten* **222**, 21 (1924).

Carl Wilhelm Wirtz (1876 – 1939), the astronomer, should not to be confused with Carl Eugen Julius Wirtz, (1910 – 1994), the nuclear physicist who worked with Werner Heisenberg.

- Additionally, Edwin Hubble did not discover that the redshift was increasing with distance;
- "Es bleibt also schon so kein Zweifel, daß die positive Radialbewegung der Spiralnebel mit
- Translation: "So there remains no doubt that the positive radial motion [i.e., *redshift* interpreted as a

#### 24

Mt. Wilson Contr. Nr. 132 (1917), Nr. 186 (1920) entlehnt. Den folgenden Ergebnissen einiger kleiner Rechnungen liegen die großen Achsen der Nebel in Bogenminuten zugrunde, aus naheliegenden Gründen nicht die Numeri, sondern die log Dm. Durch Zusammenfassung in geeignete Gruppen nach den beiden Argumenten log Dm und v erhält man das Täfelchen:

Argum. log Dm			Argum. $v$		
$\log Dm$	v	n	υ	log Dm /	n
0.24	+827 km	9	— 183 km	1.11	5
0.43	+656	7	+ 255	0.76	8
0.66	+512	8	+475	0.96	6
o.88	+555	10	+ 630	0.52	8
1.07	+334	5	+ 809	0.48	7
1.71	- 20	3	+1180	0.60	8

Click for document.



#### **Carl Wilhelm Wirtz**




The title, in French, may be translated into familiar colloquial English as "same difference" or "history repeats itself." However, now is the time to change history, to boldly stand up and say "No more of this <u>fraudulence</u>!"

The replication crisis ..., in which "sexy" papers fail to stand up to closer scrutiny, can be blamed in part on scientists being motivated by a need for attention and authority as well as curiosity about the natural world... – Bruno Lemaitre

Quoted from: Hannah Devlin, "Science falling victim to 'crisis of narcissism'," The Guardian (20 Jan 2017).

Plus ça change, plus c'est la même chose.



An Essay on Science



### The Need for Freedom in Scientific Research (original essay title)


The history of science teaches that the greatest advances in the scientific domain have been achieved by bold thinkers who perceived new and fruitful approaches that others failed to notice. If one had taken the ideas of these scientific geniuses who have been the promoters of modern science and submitted them to committees of specialists, there is no doubt that the latter would have viewed them as extravagant and would have discarded them for the very reason of their originality and profundity. As a matter of fact, the battles waged, for example by Fresnel and by Pasteur suffice to prove that some of these pioneers ran into a lack of understanding from the side of eminent scholars which they had to fight with vigor before emerging as the winners. More recently, in the domain of theoretical physics, of which I can speak with knowledge, the magnificent novel conceptions of Lorentz and Planck, and particularly Einstein also clashed with the incomprehension of eminent scientists. The new ideas here triumphed; but, in proportion as the organization of research becomes more rigid, the danger increases that new and fruitful ideas will be unable to develop freely.

Let us state in a few words the conclusion to be drawn from the foregoing. While, by the very force of circumstances, research and teaching are weighted down by administrative structures and financial concerns and by the heavy armature of strict regulations and planning, it becomes more indispensable than ever to preserve the freedom of scientific research and the freedom of initiative for the original investigators, because these freedoms have always been and will always remain the most fertile sources for the grand progress of science.

Louis de Broglie, April 25, 1978

Louis de Broglie, Heisenberg's Uncertainties and the Probabilistic Interpretation of Wave Mechanics, (Kluwer, Dordrecht, 1990), p. xviii.





For many centuries, astronomers and mathematicians worked on the details of the geocentric cosmology of the Ptolemaic system, which standardized the Geocentric model in the academic world of the second-century CE. Many lifetimes and man-centuries of effort were devoted to this futile effort, as was also the case with alchemy, commonly associated with the "Dark Ages."

Currently, there are dozens of programs, typically involving enormous public investment and the careers of hundreds of trained scientists and support personnel, investigating 'dark energy' and 'dark matter,' which are equally futile. Although not certain, prior to required prediction verification, it is likely that "gravitational wave" research, according to the current conventional approach, is similarly futile. In modern times, the Nobel Prize is more than an award for personal achievement; it establishes a line of academic research as requiring decades of effort by the global scientific community, effectively standardizing the awarded work, which then garners the better part of human and capital resources. Accordingly, when such a prize standardizes a futile avenue of research, this inevitably proves to be catastrophic, both for nascent scientific careers and for the advancement of human knowledge; time and resources may be misdirected for many decades.

The predictive equations put forward in this report were first put forward in 2005, when the author was a "Visiting Scholar" at Stanford University. Between 2005 and 2010, most of this report's content was communicated to authorities at leading academic institutions worldwide (e.g., this 2010 report). The response included contrived ad hominem attack (prior to MG12) and, the misleading awards of the 2006 (2005+1) and 2011 (2010+1) Nobel Prizes in Physics.







# Detailed online abstract from MG12 (July 2009)

### **COM2** - Dark Energy and Universe Acceleration Speaker Mayer, Alexander Talk Title Cosmological Implications of the SDSS and 2dF Redshift–Population Histograms Abstract Empirical observation of galaxy population density in redshift space is inconsistent with expectations based on the canonical cosmological model. The magnitude of this inconsistency implies a significant error in the canonical model's theoretical redshiftdistance relationship. A new model has been derived that rests on first principles and which is consistent with observed galaxy population density in redshift space. Additionally, the observed Type Ia supernovae redshift-luminosity curve does not require interpretation as accelerating cosmic expansion. Moreover, the revised redshiftdistance relationship of the proposed new model implies that the entire mass of earlytype galaxies associated with observed Einstein rings is composed of normal matter; no "dark matter" need be assumed to exist in these systems. Talk view

This abstract should be viewable online under L. Cosmological Models COM2: A







# Geocentrism and Big-Bang cosmology

This is an illustration of the Ptolemaic model of the solar system, which involved tremendous intellectual effort by many people over generations and much complex mathematics. That did not prevent the fact that it is *utter nonsense* due to a fatal false premise.

In light of this report, the Big-Bang theory is similar *utter nonsense*. The Universe is not expanding from a singularity, which is a figment of the imagination that has no correlation to physical reality. The idea that the entire mass-energy of the Universe (some trillion galaxies) was, just some billions of years ago, compacted into an infinitely small "Cosmic Egg" is truly *idiotic*; no other term adequately applies to that bizarre, irrational idea, concocted by a religiously-motivated young Catholic priest, likely suffering from severe post-war trauma.

At least this absurdly complex model could make quite accurate ephemeris predictions, so it was not completely *idiotic*. The same Big-Bang cosmology is *pseudointellectual garbage*; it is high time to cannot be said for the Big-Bang theory, abandon that ill-conceived theory. A failure to do so in the face of which has no correlation to empirical reality, this new information is an indication of mental and moral disorder. and arguably encourages irrational thinking.

